Important topics Anatomy - by DR. S.D

- 1. UPPER LIMB IMPORTANTS -
 - BRACHIAL PLEXUS
 - AXILLA
 - AXILLARY ARTERY
 - CUBITAL FOSSA
 - MAMMARY GLAND
 - MAMMARY GLAND LYMPHATICS
 - RADIAL NERVE
 - ULNAR NERVE
 - MEDIAN NERVE
 - ROTATOR CUFF MUSCLES
 - FLEXOR RETINACULUM
 - MID PALMAR SPACE
 - SNUFF BOX
 - LUMBRICALS
 - THENAR
 - SHOULDER JOINT
 - ELBOW JOINT
 - SUPINATION & PRONATION
 - CARPEL & META CARPLES
 - IMPORTANT CLINICAL ANATOMY (REFER BDC)

2. LOWER LIMBS IMPORTANTS -

- FEMORAL TRIANGLE
- FEMORAL SHEAT & CANAL
- FEMORAL ARTERY
- FEMORAL NERVE
- ADDUCTOR CANAL
- POPLITEAL FOSSA
- OBTURATOR NERVE
- SCIATIC NERVE
- ARCHES OF FOOT
- LOWER LIMB VENOUS DRAINAGE
- HAMSTRING
- FLEXOR RETINACULUM
- HIP JOINT
- LOCKING & UNLOCKING OF KNEE
- KNEE JOINT
- IMPORTANT CLINICAL ANATOMY (REFER BDC)

3. THORAX IMPORTANTS -

- PLEURA
- RECESSES OF PLEURA
- PULMONARY LIGAMENTS
- FISSURE & LOBES OF LUNGS
- ROOT OF LUNG
- BRONCHO PULMONARY SEGMENTS
- MEDIASTINUM
- PERICARDIUM
- RIGHT ATRIUM
- HEART BLOOD SUPPLY
- VEINS OF HEART
- AZYGOS VEINS
- ARCH OF AROTA
- THORACIC DUCT
- INTERCOSTAL NERVE
- INTERCOSTAL SPACE
- INLET OF THORAX
- OUTLET OF THORAX
- IMP CLINICAL ANATOMY (REFER BDC)

ANATOMY OF CECUM

M56B00KSTORE - WRITTEN BY - DR.SAHITHI (PHYSIOTHERAPIST)

INTRODUCTION

- The cecum is blind end /cul-de-sac of the large intestine below the level of ileal opening.
- It projects into the right iliac fossa.
- Cecum lies above the lateral part of inguinal ligament.

DIMENSION

- 6cm wide
- 8cm long
- Often mobile
- Enveloped by peritoneum

EXTERNAL FEATURES

- 1. TENIAE COLI
 - There are three ribbon like bands of the longitudinal muscle coat.
- 2. SACCULATION
 - These are a series of pouches /dilations in the wall of cecum and colon.
 - The sacculation are responsible for the characteristic puckered appearance of the large intestine
- 3. APPENDICES EPIPLOICAE
 - These are small bags of visceral peritoneum filled with fall attached to the teniae of large intestine.

TYPES OF CECUM M56BOOKSTORE

On the basis of growth of the cecum four types of caecum may be found in adults:

- **CONICAL TYPE /FETAL TYPE:** 2% caecum is conical shape.
- **INFANTILE:** The caecum is quadrate in shape.
- **NORMAL TYPE**: 80-90% the right saccule is larger than the left.
- **EXAGGERATED TYPE**: 4-5% the right saccule is hugely enlarged and the left saccule is absent/ atrophiles.

PERITONEAL REALTIONS

- The about 90% individual the cecum is almost completely surrounded by the peritoneum.
- The vermiform appendix frequently lies in the retrocaecal recess.

VISCERAL REALTIONS

- 1. ANTERIOR:
 - Coils of intestine
 - Anterior abdominal wall
- POSTERIOR:
 - MUSCLES: Right psoas iliacus.
 - VESSELS: Testicular or ovarian.
 - NERVE: Genitofemoral femoral and lateral cutaneous nerve of thigh.
 - Appendix in the retrocaecal recess.

M56BOOKSTORE

POSTERIOR REALTIONS

- 1. Genito femoral.
- 2. Femoral nerve.
- 3. Lateral cutaneous nerve of thigh.

ILEO CECAL JUNCTION

The junction of cecum with the ascending colon demarcated by the entrance of the ileum.

POSSIBLE FUNCTION OF THE ILEO CECAL JUNCTION:

- Preventing reflux from the cecum to the ileum.
- Regulating the passage of contents from the ileum to the cecum.

APPENDICULAR ORIFICE

- The appendix is attached to the postero medial wall of the cecum.
- Appendix orifice lies 2 cms inferior to the ileo cecal orifice.
- The orifice is guarded by insignificant semicircular fold of mucous membrane called "Valve of Gerlach" attached to the lower margin of the opening.

BLOOD SUPPLY

Branches of ilieo colic artery from the superior mescentric artery.

- The anterior cecal artery.
- The posterior cecal artery.

VENOUS DRAINAGE

• Ilieo colic vein to superior mesenteric vein to portal vein.

LYMPHATIC DRAINAGE


• The lymph vessels from the caecum drain into the ileo colic lymph nodes which in turn drain into the superior mesenteric group of pre- aortic lymph nodes.


NERVE SUPPLY:

- Sympathetic nerve supply: T11&L1 spinal segments through superior mesenteric plexus.
- Parasympathetic nerve supply: Vagus nerve.

CLINICAL ANATOMY:

- Abnormal position.
- Inflammation.
- Intestinal tuberculosis.





JOIN US ON TELEGRAM - M56BOOKSTORE

ANATOMY OF INGUINAL CANAL -

WRITTEN BY - ISHA SINGH (BPT)

Oblique intermuscular passage in the lower part of anterior abdominal wall

SITUATED: just above the medial half of inguinal ligament

- directed downward, forward, medial

EXTENDED: From deep inguinal ring to superficial inguinal ligament

DEEP INGUINAL RING:

- oval opening in fascia transversalis
- situated 1.2 cm above the midinguinal point
- immediately lateral to stem of inferior epigastric artery

SUPERFICIAL INGUINAL RING:

- triangular opening in external oblique aponeurosis
- it is like an obtuse angle triangle
- ❖ BASE from by public crest
- ❖ To margin of triangular ...lateral and medial reflected as crura
- ❖ Beyond the apex of triangle: 2 crura United by innercrural fibres

M56BOOKSTORE

BOUNDARIES:

#ANTERIOR WALL: whole extend

- skin by superficial fascia
- external oblique aponeurosis
- ❖ lateral 1/3rd internal oblique muscle

#POSTERIOR WALL: whole extend

- fascia transversalis
- extra peritoneal tissue
- parietal peritoneum

➤ MEDIAL 2/3RD:

- conjoint tendon
- medial end by the reflected part of inguinal ligament

➤ ROOF: M56BOOKSTORE

- Arched fiber
- Internal oblique
- Transversalis abdominis muscle

> FLOOR:

- Grooved upper surface of inguinal ligament
- Medial end by lacunar ligament

STRUCTURE PASSING THROUGH THE CANAL

1. spermatic cord in males


Round ligament of the uterus in female

2. Ilioinguinal nerve enters the canal through the interval between the external and internal oblique muscle and passes out through the superficial inguinal ring

INGUINAL HERNIA

> Types of hernia

- Epigastric hernia
- Incisional hernia
- Umbilical hernia
- Femoral hernia
- Inguinal hernia




PARTS OF HERNIA


M56BOOKSTORE

> 4 Parts


- Neck
- content is a loop of intestine
- Sac of hernia
- coverings of hernia




DIRECT INGUINAL HERNIA (posterior)



INDIRECT INGUINAL HERNIA



HASSELBACH'S TRIANGLE



HERNIA M56BOOKSTORE

> EXTERNAL HERNIA

- Inguinal 80%
- Incisonal 10%
- Femoral 5%
- Umbilical 4%
- Epigastric < 1%
- Others < 1%

Inguinal and femoral hernia or know as groin hernia

> SINGNS AND SYMPTOMS

- Outside lamp
- Heavy discomfort around gut
- Pain/ache on exertion
- Constipation

> RISK FACTORY

- History of hernia
- Older age
- Male sex
- Causation
- Chronic cough
- Chronic constipation
- Abdominal wall
- Smoking
 - Hernia is protrusion of any abdomina content through any of its wall. this is called External hernia
 - Intestine protrubes into the no entry zone within abdominal cavity itself this condition is called Internal hernia

TYPES OF HERNIA

- Internal hernia
- External hernia

INGUINAL HERNIA: protusion of loop of intestine through intestinal wall is called inguinal hernia

INDIRECT INGUINAL HERNIA: when the protusion occur through the deep inguinal ring , inguinal canal , superficial inguinal ring into scrotum it is called Indirect hernia

COVERINGS -

- Extraperitoneal tissue
- Internal spermatic fascia
- Cremastric fascia
- External spermatic fascia
- Skin

#DIRECT HERNIA -

• When the protusion occur through weak posterior wall of inguinal canal or traingle of Hesselbach this is called direct hernia

#HESSEIBACH -

- Bounded by inf epigastric artery, lateral border of rectus abdominis and inguinal ligament
- This area is divided into medial and lateral part by passage of obliterated umbilical artery

> COVERING

#COVERINGS OF LATERAL DIRECT INGUINAL HERNIA

- Extraperitoneal tissue
- Fascia transversalis
- Cremasteric fascia
- External spermatic fascia
- Skin

#COVERINGS OF MEDIAL DIRECT INGUINAL HERNIA

- Extraperitoneal tissue
- Fascia transversalis
- Conjoint tendon
- External spermatic fascia
- Skin

Anatomy of Vermiform Appendix

M56BOOKSTORE - Written by - Dr. Sakshi (physiotherapist)

- **POSITION:** Appendix 2cm from Iliocaecal valve
 - position is medially and length is about 2-20cm
- ORIGIN:
- -Arises from Posteromedial wall of cecum
- -Above the 2cm ilo caecal junction
 - **DIMENSIONS:**

Length – 2 to 20cm (Average 9cm)

- -It is larger in children than adults
- -Varies according to the age

Width -5mm

- -It's a tube like structure
- -Diameter of lumes varies with age

Parts of Appendix

- Base
- Body
- Tip

Body -

- -Body is tube like structure
- -Situated between Base and Tip
- -Body is narrow

Tip-

-Least vascular distal blind end



• Position Of Appendix

M56bookstore

- -Lies in right iliac fossa
- -Base of appendix is fixed

1. Paracolic Position (11'o Clock):

- -Here tip ascends right side of ascending colon
- -seen in about 2% cases

2. Retro caecal position (12'o Clock):

- -Commenest type
- -Seen in > 60% cases
- -Passes through behind the caecum and Ascending colon

3. Splenic Position (2'o Clock):

- -Passes from the pre ileal position and medially passes through ileum
- -Pre ileal variety is most dangerous (inflammation from appendix spreads to general peritoneal cavity)
- -Pre ileal seen in 1% cases
- -Post ileal seen in 0.4% cases

4.Promonteric Position:

- -Rare in occurance
- -Appendix tip is directed horizontally towards Sacral promontary

5.Pelvic position:

-Second commonest position

- -seen in >30% cases
- -Appendix tip passes downwards and medially and crosses pelvic brim to enter the pelvic

6.Mid Inguinal (6'0 clock) (Subcecal Position):

- -Appendix tip passes vertically downwards below the cecum and pointed towards the inguinal ligament
- -Seen in 2% cases M56bookstore


PERITONEAL RELATIONS

- -Vermiform appendix is intraperitoneal structure
- -Suspended by small triangular peritoneal fold
- -Derived from posterior / left layer of mesentry of ileum
- -Appendicular artery runs through the mesoappendix portions


• Appendicular Orifice

-Sometimes guarded by indistinct seminular fold of mucous membrane known as **Valve Of Gerlach**





INTERIOR OF CAECUM



Axilla - dr s.d notes

Also called cervico axillary canal

It is a pyramidal space

Present b/w upper thoracic wall & arm


Boundaries -

- 1.Apex
- 2.Base
- 3. Anterior wall
- 4. Posterior wall
- 5. Medial wall
- 6. Lateral wall

Apex is blunt – directed upwards and medially into root of neck (cervico-axillary canal)

Base – directed downwards

Diagram –



Figs 4.1a-c: (a) Dissection of axilla, (b) axilla in living body, and (c) relations to the skeleton

Boundaries description -

- 1. Apex blunt
 - Directed upwards
 - Cervico- axillary canal

Bounded by -

anteriorly - clavicle

Posteriorly – scapula & coracoid process

Medially - 1st rib (outer border)

2. Base – directed downwards

At this chest and narrow at the arm

Formed by – skin

Superficial fascia

Axillary fascia

It is convex upwards, forming concavity of armpit.

Anterior wall – formed by

- pectoralis major
- clavi pectoral fascia
- pectoralis minor

posterior wall – formed by


- sub scapularis
- teres minor
- lattismus dorsi


medial wall - formed by

- Upper part of shaft of humerus
- Coracobrachialis
- Short head of biceps brachii

Contents -

- Axillary artery & its branches
- Axillary vein
- Axillary lymph node
- Long thoracic nerve
- Intercoastal nerve
- Axillary tail of breast Brachial plexus








Fig. 4.11: The axillary lymph nodes

Reference – anatomy bdc 9th edition page no – 53

AZYGOS VEIN

THE AZYGOS VEIN MEANS – UNPAIR

FLOWCHART - AZYGOS VEIN

DRAINS INTO

THRORACIC WALL &

UPPER LUMBAR REGION

FORMS IMPORTANT CHANNEL

CONNECTING I.V.C p

- Veins occupies upper part of posterior wall of abdomen & mediastinum
- Main contents –
- 1. Formation
- 2. Course
- 3. Relations
- 4. Tributaries
- Formations –
- Union of lumbar azygos vein
- Right subcoastal region
- Ascending lumbar vein

Course –

- Enters thorax & passing through aortic opening of diaphragm
- Ascends up 4th thoracic vertebrae
- & finally pierces into pericardium through svc (superior vena cava)

Relations -

- Anterior
- Posterior
- Right
- Left
- Anterior oesophagus
- Posterior right post. Intercoastal arteries
- Right right lung pleura splanchnic nerve
- Left left lung
- Aorta
- Thoracic duct
- Trachea
- Vagus


Tributaries -

- Right superior intercoastal vein
- 5th to 11th intercoastal vein
- Hemiazygos vein (accessory)
- Right brochial vein
- Oesophageal, mediastinal, pericardial veins

- HEMI AZYGOS VEIN

Formed formly from azygos vein only -

- Also called inferior hemiazygos vein
- Mirror image of lower part of azygos vein
- Formation -
- Left lumbar
- Left ascending lumbar region
- Left subcoastal vein
- Tributaries -
- 9th to 11th (post. Intercoastal vein)
- Oesophageal vein




Azygos diagram — Refrence- B.D CHAURASIA — VOL 01 — PAGE NO — 254 9^{TH} EDITION

BRONCHIL TREE

S.D NOTES

Location – at the level of lower border of 4th thoracic vertebrae.





Right principle bronchus left principle bronchus

1. Right principle bronchus -

2.5 cm long

Shorter

Wider

More lined with trachea

M56BOOKSTORE

Infections are more common.

Making 25 degree with tracheal birfurcation

2. Left principle bronchus –


05 cm long

Longer

Narrower

More oblique

Making 45 degree with tracheal bifurcation



M56BOOKSTORE




Fig. 16.7: Parts of a pulmonary unit

Flow chart — M56BOOKSTORE



Enters the lungs through hilum

Divides into secondary lobar bronchii

There are 03 lobar bronchii on right side

02nd on the left side

Each lobar bronchus divides into tertiary bronchii

Tertiary also called segmental bronchii



1. Right side segments 2. Left side segments

10 segments

09 segments

Splits into respiratory bronchioles

Respiratory bronchiole of lung known as pulmonary unit.


Microscopic passage –

- 1. Alveolar ducts
- 2. Atria
- 3. Air sacules
- 4. Pulmonary alveoli (small cavity)

M56BOOKSTORE

Bronchopulmonary segments table -

Table 16.3: The b	ronchopulmonary segments
Lobes A. Upper B. Middle C. Lower	Right lung Segments 1. Apical 2. Posterior 3. Anterior 4. Lateral 5. Medial 6. Superior 7. Medial basal 8. Anterior basal 9. Lateral basal
A. Upper• Upper division• Lower divisionB. Lower	10. Posterior basal Left lung 1. Apical 2. Posterior 3. Anterior 4. Superior lingular 5. Inferior lingular 6. Superior 7. Medial basal 8. Anterior basal 9. Lateral basal 10. Posterior basal



CAVERNOUS SINUS

Cavernous sinus- paired dural venous sinus

Located- Within the cranial cavity

Divided by septae into two small caves

M56BOOKSTORE

Anatomical Location:

- Located within the middle cranial fossa
- Either side of sella tursica of sphenoid bone
- Enclosed by the endosteal and meningeal layers of dura mater

Borders:

- Anterior Border- Superior orbital fissure
- <u>Posterior Border</u>- Petrous part of temporal bone
- Medial border- Body of sphenoid bone
- <u>Lateral Border</u>- Meningeal layer of duramater

Roof:

- Formed by meningeal layers of the duramater
- attaches to clinoid process of sphenoid bone

Floor:

- Formed by endosteal layer of duramater
- Greater wing of sphenoid bone

M56BOOKSTORE

Contents:

Structures passing through the C.S-

- enters into orbit
- Travels through sinus
- Travels through lateral wall of sinus

Structures passing through the sinus-

- Internal Carotid artery
- Abducens nerve

Structures passing through lateral wall-

(MOTTO OK)

- i) Occulomotor Nerve O
- ii) Trochlear Nerve T
- iii) Opthalmic Nerve O
- iv) Maxillary Nerve. M
- v) Trigeminal Ganglion T

M56BOOKSTORE

Incoming channels-

- Superior Opthalmic Vein
- Inferior Opthalmic Vein
- Central Vein of Retina

Communications-

- Transverse sinus- drain through superior petrosal nerve
- Internal jugular vein- Inferior petrosal nerve
- Pterygoid Plexus- Emissary vein
- Facial vein- Sup. Opthalmic Vein
- Rt. and Left Cavernous sinus- Ant Post. Intercavernous sinus

M56BOOKSTORE

Applied Anatomy-

- Cavernous sinus thrombosis
- Clot formation within sinus


Paired Venous sinuses-

- Cavernous sinus
- Superior petrosal Sinus
- Inferior Petrosal sinus
- Transverse sinus
- Sphenoparietal sinus
- Petrosquamous sinus
- Middle Meningeal sinus


M56BOOKSTORE


Unpaired venous sinuses-


- Superior saggiftal sinus
- Inferior sagittal sinus
- Straight sinus
- Occipital sinus
- Ant. Inter Cavernous sinus
- Post. Inter cavernous sinus



Basilar plexus of veins







1. Olfactory nerve – sensory type

Location - Fore brain

Distribution - Nerve of smell

2. Optic nerve - sensory type

Location - forebrain

Distribution - Nerve of vision

SDNOTES

3. Oculomotor - Motor type

Location - Midbrain

Distribution - eye muscles

except- superior oblique & lateral rectus.

4. Trochlear - Motor

Location - Midbrain

Distribution - superior oblique

5. Trigrminal - Both sensory/motor

Location - pons vrolli

Distribution - 03 branches,

opthalamic, Maxillay and mandibular

6. Abducent – Motor type

Location - pons vrolli

Distribution - to lateral rectus

7. Facial - both - motor/sensory

Location - pons vrolli

Distribution - motor- lingual gland

Sensory- Ant. 2/3rd of tongue &

Soft palate

SDNOTES

8. Auditory - sensory

Location - pons vrolli

Distribution - nerve of hearing

9. Glosso pharyngeal – both s/m

Location - medulla oblongata

Distribution - motor - pharyngeal muscle

Parotid gland

Sensory – post. 1/3rd of

Tongue , tonsils , pharynx

Carotis sinus & body

10. Vagus - both

Location - medulla oblongata

Dist . — thorax & abdominal viscera.

11. Accessory nerve – Motor

Location - Motor

Distribution - sterno mastoid, trapezius

Constrictor muscles

12. Hypoglossal - Motor

Location - medulla oblongata

Distribution - muscles of neck & tongue

Trick -

one of our trained teacher asked for a good vehicle and horse.

SD NOTES

KIDNEY (EXCRETORY SYSTEM/URINARY SYSTEM)

M56BOOKSTORE ANATOMY NOTES

- Reddish brown s
- Bean shaped excretory structure
- Situated between T12 and L3 vertebra


Each kidney of an adult measures

- 10-12 cm length
- 5-7 cm width
- 2-3 cm thickness
- 120- 170 gm weight

Outer layer of the kidney is a tough capsule

The kidney is divided into two zones

- 1. Outer cortex
- 2. Inner medulla



Kidney Anatomy

EXTERNAL FEATURES

- Shape
- 2 Pole Upper & lower pole
- 2 border Medial border & Lateral border
- 2 surface
- 2 Hilum (anterior posterior)

INTERNAL FEATURES

- Outer Cortex
- Inner Medulla
- Renal Sinus

External Features

- Shape
- Poles

The pole is further divided into 2 parts

- 1. Upper
- 2. Lower

The upper pole is covered by supra renal gland whereas the lower pole is pointed.

• 2 borders are divided in Medial and Lateral

MEDIAL BORDER CONCAVE

• Middle part is more concave which is also known as hilum.

Hilum is further divided into 3 subparts

- 1. Renal Vein
- 2. Renal Artery
- 3. Renal Pelvis
- LATERAL BORDER CONVEX.

The lateral border is divided into 2 surfaces

- 1. anterior
- 2. posterior

Anterior surface - irregular

Posterior – flute – Abdominal Wall

Internal Features

1. Outer Cortex

- Medulla consists about conical masses called pyramid
- Outer portion of kidney
- Between renal capsule and renal medulla


2. Inner Medulla

 Medulla is divided into few conical masses called medullary_pyramids.

3. Renal Sinus

- Renal Artery
- Renal vein
- Renal pelvis
 Renal Pelvis has 2 major calyx- 7 13 Minor calyx

Kidney



ANTERIER RELATION

RIGHT KINDEY	LEFT KIDNEY
1. parts of supra renal gland	1. parts of supra renal gland
2. 2 nd part of duodenum	2. spleen
3. hepatic flexure	3. stomach
4. small intestine	4. pancreas
5. right lobe of liver	5. splenic flexure

POSTERIER RELATION


RIGHT KIDNEY	LEFT KIDNEY
1. Diaphragm	1. Psoas major muscle
2. Accurate ligament	2. Quadratus lumborum
3.Tranverse abdominis muscle	3. Subcostal vessels

Right kidney

- · Right suprarenal gland
- · Right lobe of liver
- · 2nd part of duodenum
- · Right colic flexure
- · Coils of jejunum

Left kidney

- · Left suprarenal gland
- Stomach
- Spleen
- Pancreas
- · Left colic flexure
- · Coils of jejunum



Kidney consists of:

- Medullary Pyramids
- Columns of Bernini
- Hilum
- Renal pelvis
- Nephron


SECRETORY PART

- Nephron
- Bowman's Capsule
- Collecting Duct

NEPHRON

BOWMAN CAPSULE	RENAL TUBULE
1. Afferent arteriol	1. PCT [PROXIMAL CONVOLUTED TUBULE]
2. Efferent arteriol	2. DCT [DISTAL CONVOLUTED TUBULE]
	3. HENLE'S LOOP
	4. COLLECTING DUCT

Nephron



BLOOD SUPPLY

- Right and left renal artery
- Abdominal aorta

NERVE SUPPLY

Renal plexuses

VEIN Renal vein

URETER -

Ureters are pair of narrow thick walled muscular tubes which carry the urine from the kidneys to the urinary bladder.

Each ureter is 25 cm long and 3mm in diameter.

The ureter is divided into 3 parts:

- 1. Pelvis
- 2. Abdominal
- 3. Pelvic

PELVIS	ABDOMINAL	PELVIC
1. Funnel shaped dilation at the upper.	Pass downwards and beneath the peritoneum.	Downwards towards peritoneum along greater sciatic notch
2. found by union of calyx	Enters in pelvic cavity by common iliac artery.	Oblique part – forward and medial through supra lateral angle of bladder
3. pass downwards and through hilum		Intra vesicle part – undergoes oblique and downwards through the bladder.
4. capacity – 5-7 ml		

Ureter starts with the renal sinus.

Descends along medical margin then downwards

POSTERIER RELATION - psoas major and femoral nerve

TIPS OF TRANSVERSE PROCESS

ANTERIOR RELATION


RIGHT SIDE	LEFT SIDE
1. Parietal peritoneum	Apex
2. 2 nd and 3 rd of duodenum	Parietal peritoneum
3. Right colic and ilocolic vessel	2 nd and 3 rd of duodenum
	Left colic and ilocolic vessel
	Terminal part of ileum
	Sigmoid colon

BLOOD SUPPLY


- Renal Artery
- Branch of Aorta
- Branch of Vesicle

N.S.

- Sympathetic T10 L1
- Para sympathetic S2 S4




So, this is about ureter which comes under the urinary system or excretory system.



URETHRA

MALE URETHRA:

- The male urethra connects the urinary bladder to the penis.
- It is the membranous canal, conveys urine.
- Its length is 18 -20 cm.
- Extends from internal urethral orifices to external urethral orifice
- Internal Urethral Orifice Neck of the bladder
- External Urethral Orifice Tip of the penis
- The prostatic part passes through prostate about 3cm long.
- Membranous part Sphincter urethra



- Penile part passes through bulk.
- Corpus Spongiosum of penis 15 cm long
- The internal sphincter is involuntary in

INTERNAL SPHINCTER

- Supplied by sympathetic nerve.
- Sympathetic nerve controls the neck of the bladder and prostatic urethra.

EXTERNAL SPHINCTHER

- Urethra is voluntary in nature.
- Supplied by prudential nerve.
- Controls the membranous sphincter.


S

BLOOOD SUPPLY

- Branches of vesicle.
- Pudendal artery.

FEMALE URETHRA-

- Female urethra is embedded within the vaginal wall.
- Its opening is situated between the labia major and labia minor.
- The female urethra is much shorter than the male urethra.
- Female urethra is 4cm long [1.5 inches].
- Begins at the neck of the bladder
- Opens to the outside of urethra sphincter.
- The mucous of the urethra much folded and contain mucous gland.
- Group of mucous gland are known as paraurethral gland.
- These are homologous with male urethra.



URINARY BLADDER

- It is one of the 7 important parts of the human body.
- It is situated within the pelvis.
- It extends up to abdominal cavity when it is full.
- It is
- It is hollow muscular organ which acts as urine reservoir.
- Capacity 120- 32000 [cubic centimeter]

EXTERNAL FEATURES – The empty bladder is trigon [tetrahedral] in shape.

APEX	BASE	NECK	
------	------	------	--

- 1) APEX It is directed downwards
 - i) Connect to the umbilicus through umbilical ligament.
 - ii) Represents the obliterated embryonic urachus.
- 2) BASE It is directed backwards
 - i) Related to the cervix and vagina in female.
 - ii) Male base is separated from the rectum by the recto vesicle pouch.
- 3) NECK The neck is divided into 3 surfaces and 4 borders.

Surface has a superior which is further into 2 parts right and left

Border has a lateral which is divided into 2 parts of anterior and posterior each.

The lowest and the most fixed part on the prostate in male.

INTERNAL FEATURES

- Interior part of the mucosa shows irregular fold.
- Small triangular area at the base of the bladder.
- Mucous attachment of muscular cortex.
- Apex of trigon is downward and forward.

LIGAMENT: The ligament is divided into 2 parts

RIGHT LIGAMENT LEFT LIGAMENT

TRUE LIGAMENT


- Lateral and medial prostatic ligament.
- Median umbilical ligament.
- Posterior ligament.

FALSE LIGAMENT

- Medial umbilical fold.
- Medial umbilical fold.
- Lateral false ligament.
- Posterior false ligament.


BLOOD SUPPLY

- Superior, inferior vesicle arteries.
- Vaginal artery.
- Gluteal artery.
- Uterine artery.



APPLIED ANATOMY -

- Nephritis
- Renal stones
- Pyelonephritis
- Tuberculosis 0f kidney
- Renal tumor
- Renal oedema
- Hypertension
- Renal failure
- Polycystic kidney
- (Chronic renal failure) dialysis needs to be done
- Renal colic due to pain in ureteric stone
- Prostate gland enlargement
- Cystoscopy
- Urinary tract infection
- Urinary incontinence




FACIAL NERVE & BELLS PALSY

PRESENTED BY – SHAMA
(119-123) SARANSH D
SARAN RAJ
SHALINI
SANDHIYA

FACIAL NERVE CONTENTS -

- Introduction
- Formation
- Fibers
- Functions
- Nuclei
- Origin & course
- Branches & distribution



Competency based presentation -

A 40-year-old male comes to the OPD with the history of inability to close the right eye and dribbling of saliva on the right side of mouth since early morning. He gives history of exposure to cold wind while travelling on the bus. On examination, there is loss of wrinkling on the right side of forehead and angle of the mouth is deviated to the left side.

INTRODUCTION-

Facial nerve is a vital nerve for facial expression.


Facial nerve – seventh cranial nerve

Facial nerve – nerve of 2nd branchial arch/pharyngeal arch –


2nd branchial arch forms –

- Stapes
- Styloid process
- Stylohyoid ligament
- Smaller cornu of hyoid bone
- Superior half of hyoid bone

Also forms – facial nerve & stapedial artery




Formation -



Muscles of Second Pharyngeal Arch

- · Muscles of facial expression
- · Occipito frontalis
- Platysma
- Stylohyoid
- Post belly of digastric
- Stapedius
- Auricular muscles



Second Pharyngeal Arch

Nerve : Facial nerve


Artery: Stapedial artery (only during fetal life)

Fibers -

- Sensory fibers (afferent)
- Special sensory fibers
- Visceral/autonomic motor (efferent)
- Somatic motor fibers

What are fibers?

A nerve fiber is a threadlike extension of a nerve cell (the neuron) and consists of an axon and myelin sheath.



Sensory fibers –

Transmitting signals to brain -----from external acoustic meatus as well as skin over mastoid & lateral pinna.

Special sensory fibers –

Receving & transmitting taste information from anterior 2/3rd of tongue.

- Visceral/autonomic motor fibers innervating -----
- Lacrimal gland
- Submandibular gland
- Sublingual gland

then innervating mucous memb.of nasal cavity & hard – soft palate.



Allowing production of tears & saliva.

Somatic motor -

innervating muscles of facial expression-----Like -

- Muscle in scalp
- Stapedius muscle of ear
- Posterior belly of digastric muscle
- Stylohyoid muscle

Nuclei -

- Nerve fibers connected to 4 x nuclei situated in lower pons
- Motor nucleus
- Superior salivatory nucleus
- Lacrimatory nuclei
- Nucleus of tractus solitarius

NTS cranial nerves ? Imp – 2mark Q

COURSE — Brainstem ------parotid gland

- Attached to brainstem by ---- 02 roots of motor & sensory
- 02 roots of facial nerve ----- attached to lateral part of pons.

• 02 roots run laterally & forward with 08th cranial N. (VBC) to reach

Internal acoustic meatus

- In meatus 2 x roots sensory & motor fuse
- 1st part laterally above vestibule
- 2nd part medial wall above promontory
- 3rd part vertically downward behind promontory.




• Facial nerve leave skull by passing through stylomastoid foramen.

Extra cranial course -

- Facial nerve crosses ------lateral sidebase of styloid process.
- Enters posteromedial surface of parotid gland.
- Runs forward through the gland ----crossing retromandubular vein &ECA.
- Behind the neck of mandible -----divided into 5 x terminal branches emerge
- Anterior border of parotid gland.

Divided into 5 x terminal branches -

- 1. Temporal branch
- 2. Zygomatic branch
- 3. Buccal branch
- 4. Mandibular branch
- 5. Cervical branch



Branches & distributions -

With in facial canal –


- greater petrosal nerve
- > stapedius nerve
- Chordatympani

Exit from stylomastoid foramen –

- Posterior auricular
- Post. Belly of digastric
- Stylohyoid

Terminal branches – with in parotid gland –


- ➤ Temporal branch
- >Zygomatic branch
- ➤ Buccal branch
- ➤ Mandibular branch
- ➤ Cervical branch



Facial nerve (VII) Scheme/Overview Nervus facialis Responsible for providing motor innervation to facial muscles, as well as taste (ant 2/3 of tongue) and producing saliva and tears Motor nucleus Internal Medullo-Acoustic pontine Meatus Nerve of Pterygopalatine Lacrimal gland, sulcus Foramen Pterygoid Superior ganglion Pterygoid mucous glands of the lacerum salivary canal **Greater Petrosal Nerve** nasal cavity, maxillary nucleus sinus and palate Spinal nucleus of trig. nerve Nerve to the stapedius Stapedius Nuclei of solitary tract Facial Submandibular Petrotympanic Submandibular & canal ganglion fissure sublingual glands Chorda tympani ^ Ant 2/3 of tongue Occipitofrontalis Occipital branch Stylomastoid foramen Stylohyoid branch Posterior auricular nerve Auricular branch Digastric branch Intrinsic auricular Superior Branch muscles Temporal branches Inferior Branch Parotid gland Zygomatic branches skin around the external Buccal branches Parotid Plexus acoustic meatus and the Marginal mandibular branch retroauricular region Cervical branch

Bell's palsy –

- Sudden paralysis of nerve at stylomastoid foramen----result in asymmetry of corner of mouth
- Inability to close to eye
- Wrinkling of skin of forehead on the same side.




How to remember - bell's palsy -

Trick -

- > B blink reflex abnormal
- > E Earache
- **▶** L lacrimation
- > L loss of tase
- > S sudden onset
- P paralysis of facial nerve

• Lesions – bell's palsy –



- A 40-year-old male comes to the OPD with the history of inability to close the right eye anddribbling of saliva on the right side of mouth since early morning. He gives history ofexposure to cold wind while travelling on the bus. On examination, there is loss of wrinklingon the right side of forehead and angle of the mouth is deviated to the left side.
- 1. Identify the condition.
- 2. Which cranial nerve is affected?
- 3. Is this an upper motor neuron or lower motor neuron type of lesion?
- 4. Which part of the nerve is involved in the lesion?
- 5.Mention the nucleus, course, branches and distribution of involved nerve.
- Answers –
- 1. The condition is Bell's palsy.
- 2. The cranial nerve affected is the facial nerve (CN VII).
- 3. Bell's palsy is typically a lower motor neuron type of lesion.
- 4. The lesion is likely affecting the facial nerve as it exits the skull.
- 5. The facial nerve originates from the facial nucleus in the pons courses through the internal auditory canal branches out to innervate the muscles of facial expression including the forehead, eyelids, nose, and mouth.



Femoral sheath

Funnel shaped sleeve of fasica – encloses 4cm

Walls -

- 1. Anterior wall
- 2. Posterior wall
- 3. Inferior

Anterior wall – form – fascia transversalis

Poetrior wall – form – fascia iliaca


Inferior - sheath merge with connective tissues


Sheath divided into 03 compartment by septa –

- 1. Lateral compartment femoral artery
- 2. Intermediate compartment femoral vein
- 3. Medial compartment also know as femoral canal

Contents of the femoral sheath:

- 1. Femoral artery
- 2. Femoral vein
- 3. Femoral branch of genitofemoral nerve
- 4. Femoral canal, which contains Cloquet's lymph node (the most superior deep inguinal lymph nodes).





Femoral canal

Medial compartment of femoral sheath


- 1. Conical in shape
- 2. Wide in above
- 3. Narrow below

Base of femoral canal also known as femoral ring Clinical anatomy –

Femoral hernia – potential weakness in abdominal wall Abdominal content may bulge out from femoral hernia

Femoral nerve – paralysis of quadriceps femoris

Sensory loss – medial side of leg



Femoral triangle

DR S.D NOTES

It is a triangle depression – infront of upper 1/3rd of thigh

Thigh below inguinal ligament

Boundaries -

- 1. Lateral medial border of sartorius muscle
- 2. Medial medial border of adductor muscle
- 3. Base inguial ligament
- 4. Apex lateral & medial border cross
- 5. Roof skin, superficial fasica, deep fascia
- 6. Floor psoas major , ilacus lateral Adductor longus , pectineous medial


Superficial fasica content -

- 1. Superficial inguinal ligament
- 2. Genitofemoral nerve
- 3. Ilio inguinal nerve
- 4. Great saphenous vein
- 5. Femoral artery

Contents –

- > Femoral artery
- > Femorak vein
- > Femoral sheath
- > Deep inguinal ligament
- ➤ Nerve

- Femoral artery 06 branches –
- 1. 03 superficial branches
- 2. 03 deep branches
- Femoral vein femoral vein



Corresponding branch of femoral artey

- Femoral sheath encloses upper 4cm of femoral vessel
- Deep inguinal ligament lymph from superficial inguinal ligament

Deep lymphatic of lower limb

Glans penis

Clitoris


• Nerve – femoral nerve

Pectinous nerve


Gentifemoral nerve

Lateral cutaneous nerve of thigh

Dr s.d notes



FEMORAL TRIANGLE



FERTILIZATION

- (pronuclei) to form a zygote.
- · it takes place within 24 hors of ovulation.
- roccurs at the Ampullary-Isthimic junction of follopian tube.
- * Fertilization has three process -
 - Syngamy -> union of male and female gamete.
 - plasmogamy -> intermixing of cytoplasm.
 - c) Karyogamy -> fusion of pronuclei of ovum and speum.
 - of Amphimizia -> the intermingling of their chromodomes
- · The Results of fertilization are -,
 - e) determination of genetic sex of the embryo.
 - 5) restoration of diploid no of chromosomes.
- initiation of cleavage.

Approximation of gameter -,

- · During coitus, male ejaculates about 200-300 million speums into the female's vagina.
- The journey of sperms from the vagina to the Ampulla of whering tube is promoted by prostaglanding present in the semen.
- The prostaglanding came powerful contractions of the utenine mucle, which is facilitated by oxytonin vieleased by Meurotlypophysis.

- in the werine cawity.
- · Consequently the sperms are sulfed in from vaging into the whering cavity.
- propulaire action, and to some extent it is also assisted by fluid currents created by wherine cilia.
- of the fallopian tube due to citiary movements.
- The sperms again become motile due to chemoattriac -tands produced by cumulus cells surrounding the ocyte.
- Before fertilization, the sperm must gain the ability to disintegrate the various occyte barriers.

 There are
 - of First barrier -, celle of corona radiata.
 - b) second barrier _, zona pellucida (formed by glycoprotein ZP1, ZP2, and ZP3)
 - c) Third barrier -> vitelline membrane of oogte.
 - This Ability to penetrate there barriers by sperm is achieved by two processes
 (i) Capacitation (occurs in female genital Treat)
 - (ii) Acrosome reaction.


- female genital tract.
- During capacitation, the glycoprodeins coat and reminal plasma proteins covering the plasma membrane of the sperm in the acrosomal region are removed.
- and then can fertilize ovum.


Acrosome Reaction ->

- the zona pellucida.
- The Acronomal reaction is induced by the zona protein.
- The Acronomal reaction deads to the recleare of the enzymes from acrosomal cap.
- r There enzymes are Hyalwonidase and Acrosin.
- radiata.
- . Acrosin (Zona Lysin); penetrates the zona pellucida.
- The Antigenic coating of the speum initiales the immunological reaction between the oocytest Fertilizin and the speum's Antispertilizin.

```
Mechanism of Fertilization ->
           Movement of sperms towards 2° oogte
  penetration of corona radiata by Hyaluronidare
  Adherance of sperm to ZP3 viereptors on zona pellucida
  Sperm bind to receptor on ZP Leads to initiation
          of Acrosomal reaction. - Acrosin released
   Aurorin facilitates penetration of sperm through Z.P
    Fulion of Aperm and membrane of 2° oocyte
    phagocytoris of sperm by 2° oocyte.

(Fait block polyspermy)
    completion of Meionia II of 20 oogte
    Atructural changes in Z.P through confical reaction
                  coortical granules in perivitelline
     discharge of
         space forme Fertilization membrane.
                  (Slow block polyapenmy)
            Speum enter the ovum
     event 1s followed by plaimogamy, Karyogamy
                and Amphimizid
                Fertilization complete.
```





CHANGES AFTER BIRTH -

- 1. Lungs starts functioning
- 2. Umblical vein -----ligamentum teres
- 3. Ductus venosum -----ligamentum venosum
- 4. Foramen ovale closes
- 5. Ductus arteriosus -----ligamentum arteriosum
- 6. Umblical arteries -----umblical ligaments
- 7. Placenta removed & delivered

Developmental components -

- Right atrium atrial chamber proper
 Right horn of sinus venosus
 Interatrial septum
- 2. Left atrium atrial chamberAbsorption of pulmonary veins
- 3. Right ventricle bulbus cordis rough part

 Conus cordis smooth part
- 4. Left ventricle primitive ventrice rough part Conus cordis smooth part
- 5. Interatrial septum septum primum fossa ovalis Septum secundum – limbus fossa ovalis
- 6. Interventricular septum Av cushion (atrioventricular)
- 7. Truncus arteriosus forms ascending aorta & pulmonary trunk.

Trick – LAB & RAT – LEFT ATRIUM, RIGHT ATRIUM (BICUSPID & TRICUSPID)

System · (Double Cheulation) Blood circulation oxygenated Blood. Blood. Pulimonevu Biscuspid veint Tricustia value APOK circulation (reins - contain Systemic Cerculation > Arteries Contain nein- No M56BOOKSTORE

Posterior Interatedal septum Pulmonary Cocoiary RA Sterno Costal Tutomenticular Diaphragmatic surface Interneuticular sulces Surface - Pulmonery surface - storno costal surface -- storno costal surface -- storno costal surface -Surface . coronary sulus (Atriron and ventricles) seperate) - Anterior Interneuticular suleus.

Internal structure of Right atrium Superior hall Anterior Superior Rightauvicle (Atrium) Cowa Oval Fossa (Medial septum) (Informenticular Latera Enteratrial · value of Ifve Posterior wall Infariore Viena walls - Anterior - Lateral - Superfor - Postorion = - Medial -- Inferior

Internal structure of Right yentricle Tricuspid value brows Chordae & Tendrace Papillary o Pulmonary · Tricuspid value Parete value - Septer cusp - Semilunar Doublehayer - Postvior eusp of Endowarding

Internal Structure of left atraining heft atrio venttillar Medial Interatrial septum. Bienspid value mifral value · Left ventricle 3 Semilinar Posterior alsp Anterios Bicustid eft Africo renticular opening Anterior wesp. ____ - Postovior Misp - (2)

Internal Structure of left atrium heft atrio venticular Interatrial Bicuspid value mifral value · Left ventouële_ Anterior Bicuspid - Anterdor wesp. -- Postorior Msp

Veins - Deoxygenated Arteries - oxygenated ful monary arteries Superior Carry deoxygenated Folmowary - Gets deoxygenated B.V velus Blood fring carrios Prevents Back flow of 181009. Infacior vanacava Gets deoxygenated Blood from Double circulation houser Partof the Body divergenated. Lungs. Heart Bode oxygeneted oxygenated

BLOOD Vessels carries Blood) Blood capplaries veins. Asteries Deoxygenated Blood oxygunted Blood

· Relations of Heart - (Syntopy) - Anteriorly - Sternum

2-8th costal cartilages

and Intercognal spaces of Both sides. Posteriorly - Esophorgus
- descending thoracic thoracic Vertebrae - Laterally - Lungs and plewer - Inferiorly - Contral tendon of diaphragm Superiorly - Aorta
- Pulmonary frank - Superior vena cono

Serous paricardium Fibrous Pericardium Endocordim Endo cardiny -- Lined by endothelium - Inverment hayor of the Heart Mainly values covered. (2) Myocardium -- Mainly Muscles fibres. - Found fibrous rings (values)

(v) - Myseardium Attria - 1 1) Atria - 02 housers

- Superficial hayer - Circular Muscles fittres

- Ima, trainal Muscles Deep hayer - Longitudinal Muscles fibres 2 Ventracles - 03 hayers - Superficial hayer — (1) Longifudinal - Middle houyer — (2) circulare M.F. - Deep hayer — (3) hongifudinal M.F. Cirular superficial -> Apex (downward) -> Deep hongifudinal Connection -(Superficial hayer)

eficarding Usceral hayer of Serous Perscardirmy Pericardium Pariental Lamine Fibrous Pericardium Pericardial (Serous Mid) Pericarding Wiceral Serous Pericardium.

Conducting System -2) Atrûoventrular Mode eft Posterior division:

Aorta (Arises from Leftrentricle) Flowchart Idliag ham - Aosta - 03 Part -(2nd Right - Ascending a osta STERNALAngle -Arch of Gorta - Descending about a Thoracic 1) Ascending aveta _02 Abdonina abrta Common iliae Artery 1 Descending about - 03 Parts Terentral Thoronic ! Abdomina Left subclaviant artery /Rt common carofild artery Rt. Subclavian artery. -RCCA External Cartoid Watery

Tuternal Carolid artery.

Thoracic Aorta and its Branches

Aorta


Aorta

De la lordifferent types

of Brandies Desending Aosta Parsenta Wiscera Branches. Branches (4) Post inter Pericardia Branch (1) coasfal Branch Brondia Branch Subcoustal (2) Branches 3 Brand Superior Branch

Post. Intercoastal 9 paires (3rd-11th IC)
Space Right Bronchia superior phrenic autery.

Abdominal Aorta



(2) Superior Mesentric artory supplies - Doplanding duodenum = jefnum iteum Calum - Appendix - Ascending wolon. - R. franswerse

Juster Mosentir Artery supplies flinds

Topography of Heart - Holdopy - hocapion of olegan in Regions of Body.

- skeletopy - Projection of organ on skeleton

- syntaly - Position of organ According the Another

(Relations of organs. Heart) Horizonter Line Left Border
Apex
3rd to stherib 3rd Jub Right Broder 3 to 5.

(Intercoasta)

Space) 379 5th reliastinum Parastuna Vine) Lowerborder - 5th rib (5th intercoastal

Hip joint M56BOOKSTORE

Type – ball & socket variety of synovial joint.

Articulating surface - head of femur with acetabulam cavity

Hip joint – high degree of stability & mobility.

Stability depends upon -

Depth of acetabulam tension

strength of ligament (SURROUNDING OF MUSCLES)

ligament -

- 1. Fibrous capsule
- 2. Iliofemoral ligament
- 3. Pubofemoral ligament
- 4. Ischiofemoral ligament
- 5. Head of femour ligament
- 6. Acetabular labrum

Ligaments description –

- 1. Fibrous capsule attached to acetabulam labrum
 Infront intertrochanteric line
- Iliofemoral ligament strongest ligament
 Triangular in shape
 Apex is attached iliac spine
 Base intertrochentric line
- 3. Pubofemoral ligament support joint

 Triangular in shape

 Superior obturator crest

 Inferior iliofemoral ligament
- 4. Ischiofemoral ligament cover joint posteriorly
 Twisted fiber
- 5. Head of femur ligament triangular in shape

 Apex fovea capitis

 Base acetabulam notch
- 6. Acetabulam labrum fibrocartilaginous rim attached to acetabulam margin

Relations -

Anterior – femoral artery, femora nerve, femoral vein, iliopsoas

Posterior – piriformis, sciatic nerve, quadratus femoris, gluteus maximus

superior – gluteus maximus, gluteus minimus, gluteus medius

inferior – obturator externus, hamstring muscles

Blood supply – obturator artery

circumflrx femoral artery

gluteal artery

Nerve to rectus femoris


Obturator nerve

Quadratus femoris

Gluteal nerve

Movements – flexion & extension Adduction & abduction Medial & lateral rotation Circumduction

Diagram - Relations of hip joint (VOL 2 BDC - PAGE NO 163 - 9TH EDITION)



Figs 12.5a and b: Relations of the hip joint




Fig. 12.2: Fibrous capsule of the hip joint




Fig. 12.3: The iliofemoral and pubofemoral ligaments




Fig. 12.4: The ischiofemoral ligament

Applied anatomy of hip joint -

- 1. Dysplasia of hip abnormal development
- 2. Congenital dislocation femur head slips upwards
- 3. Perthes disease pseudo coxalgia
- 4. Coxa vara neck shaft angle reduced
- 5. Dislocation of hip joint
- 6. Aspiration of hip joint

Age wise paterrn -

Below 5 yrs – congenital dislocation

5 – 10 yrs - perthes disease

10-20 yrs – coxa vara

Above 40 – osteo arthritis

Dr s.d notes

Kidney

M56bookstore

- > pair of excretory oragan
- > situated on posterior abdominal wall
- retroperitoneal in the upper left & right abdominal quadrants.
- > Functional unit of excretory system.

Kidney helps in -

- > Acid- base balance
- ➤ Blood pressure
- > Homeostatic parameters

Shape & colour -

- ➤ Bean shaped
- > Reddish brown in colour

Size -

- ➤ 11 cm long
- > 06 cm broad
- > 03 cm thick
- ➤ Weight 150 gm in males

135 gm in females.

Location -

- > epigastric
- ➤ hypochondriac region
- ➤ lumbar region
- > umblical region
- ➤ Extend up to T12 to L3
- > Right kidney is slightly lower than left kidney.

External features – 222 trick

- > 02 poles
- ➤ 02 surfaces
- > 02 borders
- > 01 hilum (entrance)

02 poles -

- > Upper pole
- ➤ Lower pole
- > Broad & close with suprarenal gland

02 surfaces -


- > Anterior surface irregular
- ➤ Posterior surface flat

02 border -

- ➤ Lateral border convex
- Medial border concave middle part hilum present
- ➤ Hilum entrance part renal vein renal artery renal pelvis.

Relations of kidney -

> Anterior realtions of kidney -




Posterior relations of kidney –

Left kidney

- · 11th and 12th ribs
- Diaphragm
- · Psoas major muscle
- Quadratus lumborum muscle
- · Transversus abdominis muscle
- Subcostal vessels and nerve
- · Iliohypogastric nerve
- · Ilioinguinal nerve

Right kidney


- 12th ribs
- · Diaphragm
- Psoas major muscle
- Quadratus lumborum muscle
- · Transversus abdominis muscle
- Subcostal vessels and nerve
- · Iliohypogastric nerve
- · Ilioinguinal nerve



Coverings -

- > Renal capsule
- > Perinephric fat adipose tissue
- > Renal fascia
- ➤ Paranephric fat

➤ Renal fasica — 02 parts — fascia of gerota & zukerkandl



➤ Blood supply of kidney –

- Blood supply of kidney

- Abdominal Aorta

Renal writery

5- segmental arrivey

Lobor arrivey

Arenate arrivey

Arenate arrivey

Therlobor arrivey

Arenate arrivey

Arenate arrivey

Affirent actouble-Glomerulus.

Efferent certerible
Peritubular plexas

Interdobular yeur.

Atrenate veln.

Interdobar veln.

Lobar veln.

Lobar veln.

Penal veln.


Enfertor venacana

05 segmental branches –


- > Apical branch
- ➤ Upper branch
- ➤ Middle branch
- > Lower branch

Applied anatomy -


- > Renal agenesis
- > Renal calculus
- > Acute renal failure
- ➤ Horseshoe shape kidney
- ➤ Polycystic kidney



G. APPLIED



Horse shoe kidney



KNEE JOINT -

DR. S.D NOTES

Type – condylar synovial joint

- 02 condylar joint b/w condyle of femur & tibia
- 01 saddle joint b/w femur & patella

Ligaments -

- 1. Fibrous capsule
- 2. Ligamentum patellae
- 3. Tibial collateral ligament
- 4. Fibular collateral
- 5. Oblique popliteal ligament
- 6. Arcuate popliteal ligament
- 7. Meniscus
- 8. Cruciate ligament
- 9. Transverse ligament

Ligament description –

Fibrous capsule – Thin & deficient anteriorly

- Replace by quadriceps femoris
 Ligamentum patellae
- Femoral attachment -

Ant – deficient

Post – intercondylar line

- Tibial attachment -

Ant - tibial tuberosity

Post – intercondylar ridge

Ligamentum patellae –

- central portion of common tendon.
- Above posterior surface of patellae
- Below tibial tuberosity

Tibial collateral ligament -

- Long band of great strength
- Attachment -
- superior Medial epicondyle shaft of femur
- inferior divided into ant. & post. part
- Ant. Part tibial shaft (medial part)
- post part tibia (medial condyle)

fibular collateral –

- superior lateral epicondyle of femur
- inferior Head of fibula

oblique popliteal ligament -

- tendon of semi membranous
- attachment –
 intercondylar line & femur lateral condyle

arcuate popliteal ligament -

extend backward from fibula head to post. Border of intercondylar area of tibia .

Meniscus –

02 fibrous cartilaginous disc = lateral & medial meniscus

Cruciate ligament – very thick & strong fibrous band

- Divided into 02 -
- Ant cruciate ligament.
- Post. Cruciate ligament.

Ant cruciate — ant. Intracondylar area —— lateral condyle of femur

Post cruciate — post. Intercondylar area —— medial condyle of femur

Transverse ligament – connects – ant. End of meniscus (lateral & medial)

Relations -

Anterior – Ant. Bursae

Ligamentum patellae.

Patellar plexus of nerves

Posterior –

- 1. Middle popliteal vessels, tibial nerve
- 2. Posterolaterally gastrocnemius

Plantaris

Common perneal nerve

3. Posteromedially – gastrocnemius

Semi-tendinosus

Semi membranosus

Gracilis

Popliteus

Medially - sartorius

Gracilis

Semi tendinosus

Saphenous nerve + vein

Semi membranosus

Laterally - biceps femoris , popliteus origin

blood supply -

- 1.05 genicular branch of popliteal artery.
- 2. Descending genicular branch
- 3. Lateral circumflex artery
- 4.02 recurrent branch of ant. Tibial artery
- 5. Circumflex fibular branch of post. Tibial artery.

Nerve supply –

- 1. Femoral nerve
- 2. Sciatic nerve through genicular branch
- 3. Peroneal nerves
- 4. Obturator nerve -----post . divison

Movements -

- 1. Extension quadriceps femoris
- 2. Flexion -

biceps femoris


semi tendinous

semi membranosus


- 3. Medial rotation semi- mem. & semi tendinous.
- 4. Lateral rotation biceps femoris

Locking – vastus medialis ----- unlocking – popliteus Applied anatomy –

- 1. Osteo arthritis
- 2. Deformities of knee
- 3. Disease of knee
- 4. Injuries of menisci
- 5. Injuries to cruciate ligament
- 6. Injuries to collateral ligaments
- 7. semi membraneous bursitis
- 8. bakers cyst



Figs 12.12a to c: Sagittal section through the knee joint of right side synovial membrane (note the cruciate ligaments)



Figs 12.11a and b: Collateral ligaments: (a) Tibial; (b) Fibular






Fig. 12.13: Transverse section through the right knee joint showing the relations





Locking & unlocking of knee joint -

Mechanism that allow the knee to remain in the postion full extension without muscular effort.

Locking occur –

Medial rotation of femur -----at last stage of extension.

Diameter of lateral femoral condyle less than medial condyle

Lateral condyle articulation in extension.

Medial condyle remain unused

Locking produced by extensor muscle ----quadriceps femoris

Unlock – reverse of locking – by lateral rotation of femur Unlock caused -----by popliteus muscle.

Larynx

Larynx is the organ for production of voice or phonation Also called the sound box.

-It's air passage and act as sphincter at inlet of lower respiratory passages.

M56BOOKSTORE

Situation:

- Lie in Ant. Midline of neck
- Extending -Root of tongue to trachea
- Extend from epiglottis above to the lower border of cricoid cartilage.

In adult male: C3- C6 vertebrae

In adult female: C1-C4

#Length:

Male- 44mm Female- 36mm

Cartilage of Larynx:

- i) Unpaired cartilages
- ii) Paired cartilages

Unpaired Cartilages:

- **T** Thyroid (shield like)
- **C** Cricoid (ring like)
- E- Epiglottis (leaf like)

M56BOOKSTORE

Paired Cartilages:

- A- Arytenoid (Cup shaped)
- **C**-Corniculate(Horn shaped)
- **C**-Cuneiform (Wedge shape)

Membranes of Larynx

- i) Extrinsic ii) Intrinsic
- 1. Thyrohyoid membrane:
- Extrinsic membrane
- connects thyroid cartilage to hyoid bone
- medial and lateral parts thickened

M56BOOKSTORE

2. Fibroelastic membrane:

- intrinsic membrane
- outside mucous membrane of larynx
- divides into two parts
- above sinus: quadrate membrane
- below sinus :conus elasticus

#Constitution of larynx (Internal features)

- Cartilages
- Laryngeal joints
 - Cricothyroid
 - Cricoarytenoid
 - 1. Cricothyroid: -synovial joint
 - -permits two movements
 - the rotatory movement (transverse axis)
 - gliding movements (all directions)

M56BOOKSTORE

- 2. Cricoarytenoid: -synovial joint
 - lamina of cricoid cartilage
 - permits movements
 - i)Rotatory vertical axis ii)Gliding (all directions)

Ligaments of larynx

Extrinsic ligaments are:

- 1. Lateral thyrohyoid ligament
- 2. Median thyrohyoid ligament,
- 3. Hyoepiglottic ligament,
- 4. Cricotracheal ligament

M56BOOKSTORE

- 1. LTL originate from thyrohyoid membrane
- 2. MTL- single ligament,
 - -originates from thyrohyoid membrane
- 3. HGL- epiglottic cartilage to hyoid bone
- 4. Cricotracheal ligament, CTL Major intrinsic ligament
 - Aryepiglottic fold
 - Vestibular fold
 - Vocal fold

Muscles movement

- i) Elevation-Thyrohyoid, Mylohyoid,
- ii) Depression Sternothyroid, Sternohyoid
- iii)Opening inlet Thyroepiglottic
- iv)Closing inlet Aryepiglottic
- <u>v)Abductor</u> Cricorarytenoid(Post.)
- vi)Adductor Cricorarytenoid(Lateral)
- vii) Modulation Oblique arytenoid, Cricothyroid
- viii)Relaxation Thyroarytenoid, Vocalis

M56BOOKSTORE

Blood supply: Superior laryngeal artery/vein
Inferior laryngeal artery/vein

#Nerve supply:

(Motor) -Recurrent laryngeal nerve

-External laryngeal nerve

(Sensory) - Internal laryngeal nerve

- Recurrent laryngeal nerve

M56BOOKSTORE

#Clinical Anatomy: -Laryngeal cancer

-Laryngitis

-Spasmodic dysphonia (Vocal cord


Spasm)

#Cartilages of Lateral wall of nose


- -Sup. Nasal Cartilage
- -Inf. Nasal Cartilage
- -3 to 4 small ala cartilages

M56BOOKSTORE

A. CARTILAGE



B. INTERNAL FEATURES




M56BOOKSTORE

C. MUSCLES Vocal ligament Lamina of thyroid cartilage - Vocalis - Thyroarytenoid Anterior arch of cricoid cartilage Direction of pull/ Vocalis and -Lateral cricoarytenoid thyroarytenoid Lateral cricoarytenoid Muscular process of arytenoid Transverse-Posterior arytenoid cricoarytenoid Posterior Transverse arytenoid C cricoarytenoid

M56BOOKSTORE

D. BLOOD AND NERVE SUPPLY




M56BOOKSTORE

Ref. BDC HUMAN ANATOMY 9th Edition, VOLUME 3, Page 282

LIVER sd Anatomy notes

- It is one of the metabolic organs.
- It is the largest gland of the human body.
- Located in the diaphragm and has also occupied the right hypochondrium.
- Extend up to the left hypochondrium.
- Almost covered by the peritoneum.
- Mostly covered by the ribs and coastal cartilage.



FEATURES

M56bookstore

- WEIGHT 1-1.5 KG
- MALES 1600 GM
- FEMALES 1300 GM

SHAPE - Wedge

Surface

- 1. Anterior
- 2. Posterior
- 3. Superior
- 4. Inferior
- 5. Right

LOBES

- 1. Right
- 2. Left
- 3. Caudate
- 4. Quadrate

LIGAMENTUN VENOSUM


- Present in between the groove of IVC and tissues.
- Caudate lobe lies on the posterior surface.
- Quadrate lobe lies in the inferior surface.

PERITONEAL RELATIONS

- 1 bare area.
- Groove of IVC.

SD ANATOMY NOTES

- Gall bladder Forsa.
- Coronary ligament.




Visceral relation -

- Anterior surface related to xiphoid process
- Diaphragm on each side separate pleura & lungs
- Posterior surface is traiangular bare area right suprarenal groove for IVC
- Inferior surface gastric impression
- Right surface quadrilateral & convex
- Superior surface quadrilateral & concavity in middle

M56BOOKSTORE

Anatomy of human liver



LIGAMENTS OF THE LIVER

- There are 5 types of ligaments.
- 1. Falciform ligament.
- 2. Left ligament.
- 3. Right ligament.
- 4. Coronary ligament.
- 5. Lesser omentum.

SD ANATOMY NOTES

1. FALCIFORM LIGAMENT

• Connecting Anterosuperior to anterior abdominal wall.

2. LEFT LIGAMENT

• Superior surface to diaphragm.

3. RIGHT LIGAMENT

· Posterior surface to diaphragm.

4. CORONARY LIGAMENT.

Bare area of liver.

5. LESSER OMENTUM

· Lips of portal hepatic.

FUNTIONS

- Manufacture of bile salts.
- Emulsification of fats.
- Absorption of fats.
- Anti-coagulant heparin.
- Forms prothrobin, fibrin and aluminum.
- Contains enzymes.
- Absorbed nutrients.
- Stores glycogen, copper, iron.
- Stores vitamins A, D, E, and K.
- Activation of vitamin D.
- Released Bile Rubin, bilverdin.

BLOOD SUPPLY

- Liver received 20% blood supply.
- · Hepatic artery.
- 80% portal vein.

NERVE

- · Hepatic plexus.
- Sympathetic derived from coelic plexus
- Para Sympathetic derived from hepatic branch

Hepatic segments -

SD ANATOMY NOTES


Right lobe – divided into – anterior & posterior part


Left lobe - medial & lateral parts

Each parts divided into upper & lower part

Fissure for ligamentum teres & ligamentum venosum – make junction b/w medial & lateral segment of left lobe.

M56BOOKSTORE





Inferior surface relations of liver

• Formation of the Neural Plate:

- The process begins around the third week of embryonic development.
- The notochord induces the overlying ectoderm to thicken and form the neural plate.
- The neural plate is a flat, thickened region of ectoderm located along the dorsal midline of the embryo.

• Shaping of the Neural Plate:

- The edges of the neural plate elevate to form neural folds.
- The central region of the neural plate remains depressed, forming the neural groove.

• Elevation of the Neural Folds:

- The neural folds continue to elevate and move toward the midline.
- This movement creates a deeper neural groove.

• Fusion of the Neural Folds:

- The neural folds begin to fuse at the midline, starting in the cervical region (future neck area) and proceeding both cranially (toward the head) and caudally (toward the tail).
- The fusion of the neural folds converts the neural groove into a closed neural tube.

• Closure of the Neural Tube:

- The neural tube closes progressively from the center of the embryo outward, leaving two openings at the ends called the anterior (rostral) and posterior (caudal) neuropores.
- The anterior neuropore closes around day 25, and the posterior neuropore closes around day 27.

• Differentiation of Neural Tube:

- After closure, the neural tube differentiates into various regions that will form the brain and spinal cord.
- The cranial end of the neural tube expands and differentiates into the primary brain vesicles: the prosencephalon (forebrain), mesencephalon (midbrain), and rhombencephalon (hindbrain).
- The caudal portion of the neural tube forms the spinal cord.

• Formation of Neural Crest Cells:

- Cells at the edges of the neural folds, known as neural crest cells, separate from the neural tube and migrate to various parts of the embryo.
- Neural crest cells give rise to a variety of structures, including peripheral neurons, glial cells, melanocytes, and facial cartilage.

• Development of Surrounding Structures:

- The mesoderm surrounding the neural tube forms the vertebrae and muscles of the back.
- The surface ectoderm, which lies above the neural tube, becomes the skin.

• Establishment of the Central Nervous System:

• The closed neural tube eventually forms the central nervous system, with the brain developing from the cranial portion and the spinal cord developing from the caudal portion.

- **Primitive Streak Formation**: During the third week of embryonic development, the primitive streak forms on the surface of the epiblast, a layer of cells in the bilaminar embryonic disc.
- **Primitive Node Formation**: At the cranial end of the primitive streak, the primitive node (or Hensen's node) develops, which is essential for the formation of the notochord.
- Invagination of Epiblast Cells: Cells from the epiblast migrate toward the primitive streak, invaginate, and move inward. Some of these cells displace the hypoblast and create the endoderm.p
- **Mesoderm Formation**: Other invaginated cells move between the epiblast and the newly formed endoderm to form the intraembryonic mesoderm.
- **Notochordal Process Formation**: Cells that migrate cranially from the primitive node form a tube-like structure called the notochordal process, which extends cranially between the ectoderm and the endoderm.
- **Notochordal Plate Formation**: The floor of the notochordal process fuses with the underlying endoderm, and the cells intercalate with the endoderm to form the notochordal plate.
- **Notochord Formation**: The notochordal plate detaches from the endoderm and rolls up into a rod-like structure called the notochord. This process involves the folding of the notochordal plate into a tube, which becomes the definitive notochord.
- **Final Positioning**: The notochord lies in the midline between the ectoderm and the endoderm and extends from the primitive node to the prechordal plate, which is located cranially.
- **Signaling Role**: The notochord plays a crucial signaling role, inducing the formation of the neural tube and influencing the development of surrounding tissues, including the vertebral column.
- **Degeneration**: In humans, much of the notochord degenerates and is replaced by the vertebral column, although remnants of the notochord persist as the nucleus pulposus within the intervertebral discs.

OESOPHAGUS

M56BOOKSTORE

- It is a narrow muscular tube.
- Lies behind the trachea.
- About 23 25 cm long.
- Begins at the end of the larynges pharynx.
- Passes through the mediastinum.
- It starts from at the level of 6th cervical vertebrae
- At the level of T10.

STRUCTURE

There are 4 layers of esophagus:

MUSOCA

- Non keratinized stratified
- Lamina propla.

SUBMUCOSA

- Contains secretive glands.
- Blood vessels.

M56BOOKSTORE

MUSCULARIS

- Upper 3rd is striated.
- Middle 3rd is striated.
- Lower 3rd is smooth,

CONNECTIVE


- OUTER loose connective tissue
- Contains capillaries
- Does not produce digestive enzymes
- Secrets muscular
- Transport food to the stomach
- Called upper esophageal sphincters
- Involuntary muscular movement called peristalsis

Parts of esophagus -

- 1.cervical esophagus
- 2.thoracic esophagus

M56BOOKSTORE

3.abdominal esophagus



CONSTRICTION

- At its starting
- Cross of the aortic arch
- At the diaphragm


• Cross by the left branches

BLOOD SUPPLY

- Branches of aortic
- Branches of gastric artery
- Branches of thyroid artery

N.S.

PARASYMPASTHETIC	SYMPATHETIC
1. Laryngeal	1. Cervical ganglia
2. Esophageal	2. Thoracic ganglia






Fig. 6.10: Mechanism of deglutition (swallowing).

- Stage 1. Bolus of food made by the soft palate.
- Stage 2. Bolus lies in oropharynx; posterior nasal opening and inlet of larynx closed.
- Stage 3. Bolus has nowhere to go, it passes into laryngopharynx and esophagus.

M56BOOKSTORE

PANCREAS

- It is partly exocrine and partly endocrine gland.
- Secretes hormones like:
- Exocrine secrets pancreatic juice
- Endocrine secrets hormone like insuline
- It is a soft organ.
- 15 20 cm long.
- 2 3 cm broad.
- 1 2 cm thick.
- 90 gm. weight.
- Lies transversely posterior abdominal wall.
- Entire organ lies posterior to stomach
- Level of L1 L2 Vertebrate.

Sd anatomy notes

- Pancreas link with duodenum through duct.
- Pancreatic duct lies in posterior surface.
- Begins at the tail.
- Runs towards the neck and bends at the neck.
- Head of the pancreas direct related to bile duct.
- Ducts enter into the second part of the duodenum.
- Joints the hepatopancreatic ampulla.
- Ampulla opens in the second part of duodenum.

PENCREAS IS DIVIDED INTO 4 PARTS

HEAD NECK	BODY	TAIL
-----------	------	------

1. HEAD

- It is expanded portion.
- Lies in the cavity of duodenum.
- External features -
- 03 borders superior, inferior, right lateral
- 02 surfaces anterior & posterior
- One process uncinate process

Relations – border – superior – overlapped by 1st part of duodenum

Inferior - 3rd Part of duodenum

Right border – 3rd part of duodenum

Surface- anterior - 1st part of duodenum

Posterior – inferior vena cava

Renal vein

Right crus of diaphragm

Uncinate process - superior mesenteric vessel

2. NECK

• Slightly constricted part.

3. BODY

- Extends from the neck tail.
- It has 3 borders
- There is projection like called tuber omental.

4. TAIL

- Narrow terminal part.
- Lies at lienorenal ligament.
- It has two types of cells

	101 0 1 1110 1110
CULTURES OF EPITHELIAL CELL	ISLETS OF LANGERHANS
I CULTURES OF EFITHELIAL CELL	I ISLE I S OF LANGENHANS

- Consist of cells alpha 1 & alpha 2, f cells, beta cells, delta cells
- Secrets

Pancreatic gastrin	Serotonin	Glucagon
i anordano gadanii	00.00	o.acago

- Produce insulin.
- The exocrine part is serous gland cells acne.
- Secrets digestive enzymes called pancreatic juice.

Sd anatomy notes

FUNCTIONS - DIGESTIVE ENZYMES CONTAINS

- Trypsine
- Lipase
- Amylase
- Peptides
- Glucagon


Ducts of pancreas - 02 ducts -

- 1. Main pancreatic duct
- 2. Accessory pancreatic duct

M56bookstore

- 1. Main pancreatic duct 3mm in diameter
- 2. Lie near posterior surface of pancrease
- 3. Begins at tail, run towards right & bend at neck downwards
- 4. 02 ducts enters the wall of 2nd part of duodenum
- 5. Join and form hepato pancreatic ampulla of vater.

Arterial supply diagram -



Arterial supply -

Mainly by – pancreatic branches of splenic artery

Superior pancreatico duodenal artery

Inferior pancreatico duodenal artery

Venous suppy -

Splenic vein

Superior mesenteric vein


Portal vein

NERVE SUPPLY

- Para sympathetic Valgus nerve
- Sympathetic Splenic nerve

Important terminologies -

- 1. Ampulla of vater
- 2. Sphincter of oddi




@ MAYO FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH, ALL RIGHTS RESERVED

The ampulla of Vater is located where your bile duct and pancreatic duct join and empty into your small intestine.

Ampullary cancer forms near many other parts of the digestive system, such as the liver, pancreas and small intestine. When ampullary cancer grows, it may affect these other organs.

SPHINCTER OF ODDI -

The sphincter of Oddi is a muscular valve responsible for controlling the flow of bile and pancreatic secretions through the ampulla of Vater into the second part of the duodenum.



PAROTID GLAND

• INTRODUCTION: -Largest Salivary Gland

-Weight 25g

SITUATION: - Below external Acoustic
 Meatus

- B/w Ramus of Mandible and Sternocleido Mastoid
- -Anteriorly: overlap Masseter Muscle

M56BOOKSTORE

- CAPSULE: -Investing layer of deep cervical fascia
 - Form capsule for gland.
 - Supplied By Auricular Nerve
 - -Deep Lamina -Thin
 - -Attached-tympani Plate
 - -Portion of deep Lamina B/W
 Styloid process and mandible is
 thicker

External features

- 3 side Pyramid
- Apex-downward

M56BOOKSTORE

- 4 Surfaces : -Superior
 - -Superficial
 - -Anteromedial
 - -Posteromedial.
- 3 Borders: -Anterior
 - -Posterior
 - -Medial

Relations

- Apex
 - -Post. Belly diagastric.
 - -Cervical Branch of facial Nerve
 - -Retro mandibular vein

Surfaces:

M56BOOKSTORE

i)Superior surface.

- -Superficial Temporal vessel
- -Auriculotemporal Nerve
- -External Acoustic Meatus
- -Posterior surface of TMT

ii)Superficial surface:

- -Skin
- -Superficial fascia
- -Parotid fascia
- -Deep parotid Lymph Node M56BOOKSTORE

iii)Antero medial surface:

- -Ramus of Mandible-Masseter
- -Medial Pterygoid.
- -Lateral surface of TMT

iv)Postero medial surface:

- -Mastoid Process
- -Styloid Process.
 - Borders:
- <u>i) Anterior Border</u> Terminal Branch of facial

Nerve

- -Parotid duct
- -Transverse facial vessel

M56BOOKSTORE

ii) *Posterior border* separate: Superficial surface to posteromedial

surface

iii) *Medial border* separate: Antero Medial to posteromedial surface

Structure within parotid gland

#Arteries:

- (i) ECA-->Enter Posteromedial Surface
- (ii)Maxillary Artery leave-->

Anteromedial surface

M56BOOKSTORE

#Veins:

Retro mandibular vein --> Formed in gland with By STV + MV

- -Lower part Anterior
- -Posterior Division --> Emerge/Apex

Nerve Supply:

Branch - Temporal Branch

-Zygomatic Branch

Upper and Lower -Buccal Branch

- -Marginal Mandibular Nerve
- -Cervical Branch

M56BOOKSTORE

Clinical Anatomy:

- -Parotid Abscess
- -Mixed Parotid Tumour
- -Parotid calculi
- -Viral parotitis


Parotid duct

- -Upper 2nd Molar teeth
- -Emerges, Middle of Anterior Border
- -Pierce in Buccinator Muscle
- -Duct open- Vestibule of Mouth.

M56BOOKSTORE


Secreto Motor


- -Inferior Salivatory Nucleus
- -Glossopharyngeal Nerve
- -Tympanic Nerve
- -Lesser Petrosal Nerve
- -Otic ganglion
- -Post ganglion fibre through Auricotemporal Nerve




M56BOOKSTORE

B. RELATIONS





M56BOOKSTORE



M56BOOKSTORE

Ref. BDC HUMAN ANATOMY 9^{TH} EDITION VOLUME 3 PAGE 114

REPRODUCTIVE SYSTEM

- M56BOOKSTORE

- WRITTEN BY : DR. SWEZA MANSURI


PHYSIOTHERAPIST

CONTENTS:

MALE REPRODUCTIVE SYSTEM	FEMALE REPRODUCTIVE	
(MRS)	SYSTEM (FRS)	
- Testis	- Ovaries	
- Penis	- Vagina – i. internal	
	ii. external	

MALE REPRODUCTIVE SYSTEM


INTERNAL GENITAL ORGANS	EXTERNAL GENITAL ORGANS
- Testis	- Penis
- Epididymis	- Scrotum
- Ductus deferens	
- Ejaculatory duct	
- Prostate	
- Male urethra	
- Bulbo urethral gland	




❖TESTIS:

- 01 pair
- Primary function → release out
- 1. **LEYDIG CELLS** produce **TESTOSTERONE**
- also called **instertital cells** (present in seminiferous tubule and made up of connective tissue)
- 2. SERTOLI CELLS formation of SPERMATOGENESIS
- LOCATION: outside the body
- **TEMPERATURE**: 2-3 C
- STRUCTURE: pouch like called as scrotum

DESCENT OF TESTIS:




*** EXTERNAL STRUCTURE OF TESTIS**



POLES	BORDERS	SURFACES
- Upper	- Anterior	- Lateral
- Lower	- Posterior	- Medial

❖ INTERNAL STRUCTURE OF TESTIS:



LAYERS:

- 1. Tunica albugenia: Dense membrane of connective tissue
- 2. Tunica vascularis


SEPTA OF TESTIS :

- they are connective tissuecalled as lobule of testis
- each lobule has single tubules called as convulated seminiferous tubule
- it has sertoli cells
- once the tubule leave the lobule then it is called as **staright tubule**

EFFERENT DUCTUS

- continue with head of epididymis

*** EPIDIDYMIS:**



PARTS:

- **HEAD**: broad, consist efferent ductus
- BODY
- TAIL

- No function but serves as a reservoir of spermatozoa
- sperm cells mature and gain the capacity to move and fertilize the ovum
- sperm cells provide nutrition to epididymis

#CLINICAL ANATOMY

- orchitis: infection occurs in gonads and epidiymis head is swelled

- -Hydrocele: dehydration


#SYMTOMS:

- fever
- minute headache
- cannot sit for longer
- time period (10-20 days)

INVESTIGATION:

ultrasonography (USD)

COVERINGS AND FIXATION OF TESTIS AND EPIDIDYMIS




#TUNICA VAGINALIS:

VISCERAL LAYER

 Attached to lateral and anterior surface of testis and fused with tunica albugenia

PARIETAL LAYER

Attached with tunica albugenia



#CREMASTER MUSCLE


- Present between internal and external spermatic fascia

#SPERMATIC CORD


- formed with the help of:
- EXTERNAL SPERMATIC FASCIA
- CREMASTER MUSCLE
- INTERNAL.SPERMATIC FASCIA

#BLOOD SUPPLY

- Testicular artery
- lymph vessels
- pampiniform plexus
- testicular plexus



❖ DUCTUS DEFERENS



#04 PARTS:

- Scrotal part
- Funicular part
- Inguinal part
- Pelvic part

#JUNCTION: ampulla of ductus deferens


- ductus deferens fused with seminal glands
- forms ejaculatory duct
- enters into the prostate region called as prostatic urethra

❖ SEMINAL GLAND

- LOCATED: behind the urinary bladder above the prostate gland

#Seminal gland

- produces 50 80% of ejaculate fluid
- it has excretory duct which attaches to ductus deferens




❖ PROSTATE GLAND:


- largest male genital gland
- produces 30% of ejaculate fluid

#CLINICAL ANATOMY:


- Prostatitis
- (Enlargement of prostate gland)
- normal range 30-40 (beyond 40 then swelling will occur


SURFACES	PARTS	ZONES
- anterior	- base	- central zone
- posterior	(upwards)	 transitional
	- apex	zone
	(downside)	- peripheral
		zone






CROSS SECTION OF PROSTATE GLAND






❖ BULBO URETHRAL GLAND



- paired of accessory genital gland
- attached to male urethral duct

❖ MALE URETHRA



#03 Portions

- prostatic urethra
- membraneous urethra
- spongy urethra


PENIS:


#EXTERNAL STRUCTURE

- consist 03 pipes/ erectile bodies
- i. Copora cavernosa (02)
- ii. Corpus spongiosum (01)


(Help in ejaculation)

- unpaired erectile bodies







ROOT	BODY	GLANS PENIS
Consist :	Composed of:	- Corona glandis
- Bulb of penis	- Corpora	
 Crura of penis 	cavernosa	
	- Corpus	
	spongiosum	



CROSS SECTION OF PENIS :



GLANS OF PENIS




SCROTUM

- pouch like structure
- sac located outside the body
- Contains: testis and epididymis

#LAYERS:

- tunica vaginalis
- tunica albugenia
- tunica vascularis
- spermatic fascia
- dartos fascia



FEMALE REPRODUCTIVE SYSTEM


INTERNAL GENITAL ORGANS	EXTERNAL GENITAL ORGANS
- Ovary	- Vestibule
- Uterine tube	- Labia majora
- Uterus	- Labia minora
- Vagina	- Bulb of vestibule
	- Clitoris

OVARY

#INNER STRUCTURE


- contains oocytes with follicle
- Produces:
 - estrogen
 - progesterone

#GROWTH AND MATURATION OF OVARY:




#OVARY:

OVARIAN CORTEX	OVARIAN MEDULLA
- Outer region	- Inner region
	Contains :
	- Arteries
	- Blood vessels
	- Capillaries




*****EXTERNAL STRUCTURE OF OVARY



BORDERS	SURFACES
- Mesovarian border	- Lateral
- Free border	- Medial

• FIXATION OF OVARIES:



#LIGAMENTS:

- 1. Ovarian ligament
- 2. Suspensory ligament of ovary attached from tubal extrimity to lateral wall of pelvic

#PERITONEAL FOLD:

VISCERAL PERITONEUM	PARIETAL PERITONEUM
 Lines the wall of organs 	- Lines the internal surface of
	abdominal and pelvic wall

#LIGAMENTS OF UTERUS


BROAD LIGAMENT OF UTERUS

- (double peritoneal fold)
- It consist 03 ligament

MESOVARIUM	Fixes the ovary
MESOSALPINX	Fixes the ovarian tube
MESOMERTIUM	From the uterus

***** UTERINE TUBE / FALLOPIAN TUBE

- paired tubes
- fixed with the uterus to ovaires
- fertilization happens in uterine tube




#PARTS OF UTERINE TUBE:

1. Infundibulum: dilated portion

Ampulla : wider
 Isthmus : narrow

4. Uterine part : attached to uterus

INFUNDIBULUM




UTERUS:

- undergo cyclic transformation
- synchronized with the ovarian cycle
- this ensures that uterus is ready for implantation

#PARTS OF UTERUS

- fundus
- body
- cervix

FEMALE REPRODUCTIVE SYSTEM



#CERVIX

- supravaginal part
- vaginal part

#UTERINE CAVITY OPENING


- uterine opening of oviduct(02)
- cervical canal(01)

#CERVIX (posterior view)

- anterior lip
- posterior lip


UTERUS CONSIST 03 LAYERS

- Endometrium (inner):
 - i.stratum functionalis
 - ii.stratum basalis
- Myometirum (middle):
 - i.longitudinal (02)
 - ii.middle (01)
- Perimetrium (outer)



#PARAMETRIAL LIGAMENTS

- Cardinal ligament
- vesicouterine ligament
- pubocervical ligament
- uterosacral ligament
- rectouterine ligament



#SUPPORTING APPARATUS OF UTERUS

- perineal muscle
- levator ani muscle

***VAGINA**

- it is hollow muscular tube like sturcture

#supports:

- Rectus
- Urinary bladder
- Uterus

#ORIFICE: vaginal orifice

#WALLS OF VAGINAL CANAL:

- anterior wall
- posterior wall

#VAGINAL FORNIX


- anterior part
- posterior part

#URETHROVAGINAL SEPTUM

retrovaginal septum

(seperating urethra and vaginal canal)

*** EXTERNAL GENITAL ORGANS**



VESTIBULE

#ORIFICE:


- external urethral orifice
- vaginal orifice

#LESSER VESTIBULAR GLAND:

- responsible for lubrication


#GREATER VESTIBULAR GLAND:

- paired mucus gland
- opens in posterior part of vestibule
- secretes fluid for lubrication




***** LABIA MINORA (inner lips)

- 02 cutaneous folds
- inner surface lined by mucus



CLITORIS

- One of the most sensitive zone in FRS
- made up of corpora cavernosa
- consist glans clitoris



❖ BULB OF VESTIBULE


- LOCATED: inside of clitoris

- type of erectile tissue which is closest to clitoris

- formed by : plexus

❖ LABIA MAJORA

upper part : anterior commissurelower part : posterior commissure



❖ MONS PUBIS :

- Covered by hair at the time of puberty
- triangular in shape

RIGHT ATRIUM

Right atrium is the right upper chamber of heart.

It receives venous blood from the whole body

Pump into right ventricle through tricuspid valve

External features -

Upper right chamber elongated vertically.

Along right border of atrium ——— shallow ventrical groove which passes from ------superior vena cava to inferior vena cava is called sulcus terminalis.

It produced by internal muscular ridge called crista terminalis.

s.d

Inlet of right atrium –

- 1. Superior vena cava
- 2. Inferior vena cava
- 3. Coronary sinus
- 4. Anterior cardiac veins


Divided into 03 -----

- 1. Posterior Smooth part –Derived from right horn sinus venosus
 - Superior vena cava
 - Inferior vena cava eustachian valve
 - Coronary sinus thebesian valve
 - Venae cordis minimae
- 2. Rough anterior part Derived from primitive atrial chamber It present a series of muscular ridge called musculi pectini.
- 3. Inter arterial septum –
 Derived from primitive atrial chamber
 Annulus ovalis & fossa ovalis the prominent
 margin fossa ovalis.

Arterial supply of heart

Heart is supply by 02 coronary arteries Arising from ascending aorta

- 1. Right coronary artery
- 2. Left coronary artery



Right coronary artery

Position – smaller than left coronary artery.

Arise from anterior sinus

Course -

pass forward b/w root of pulmonary trunk & right auricle

runs downwards – right anterior coronary sulcus

winds around the inf. Border to reach – diapraghmatic surface of heart.

Terminate by anastomosis with left coronary artery.

<u>Branches</u> –

- 1. Arterial branch ant. Post. & lateral
- 2. Right conus artery called annulus of vieussens
- 3. Ventricular branch ant. Post group
- 4. Right marginal artery
- 5. Post interventicular branch

Area of distribution -

Right atrium

Great part of right ventricle

Small part of left ventricle

Conducting system of heart except left branch of av bundle.

LEFT CORONARY ARTERY -

Arises from – left aortic sinus

Course -

Run forward to left end emerge b/w pulmonary trunk

Run downward in a groove – continuation with circumflex artery

Binds left border of heart & continue with left coronary sulcus



& make anastomosis with right coronary artery.

BRANCHES & DISTRIBUTION –

- 1. Anterior interventricular branch largest
- 2. Circumflex branch is terminate part of LCA
- 3. Ant. Post ventricular branch
- 4. Arterial branch

AREA OF DISTRIBUTION -

- 1. Left atrium
- 2. Greater part of left ventricle
- 3. Ant.part of inter ventricular septum
- 4. Left branch of av bundle

VEINS OF HEART

Great cardiac vein
Middle cardiac vein
Right marginal vein
Posterior vein of left ventricle
Anterior cardiac vein
Venae cordi minimae

Coronary sinus – largest vein of heart Situated into left posterior coronary sulcus -----in right atrium.

TRIBUTARIES -

Great cardiac vein – with ant. Ventricular artery & left coronary artery

Middle cardiac vein – with – posterior interventricular artery


Small cardiac vein - with RCA

Right marginal vein – with – marginal branch of RCA Post vein of left ventricle – towards to diaphragmatic surface


Oblique vein – running of post surface of left atrium

Anterior cardiac vein – open directly into right Atrium.

HEART AND PERICARDIUM



BLOOD SUPPLY OF HEART



- It is a small, vascular organ.
- Dark purple in color.
- One of the major lymphatic organs.
- Acts as a filtration of the blood.
- Plays an important role in immune system.

SHAPE – wedge shaped.

- SIZE 1 Inch thick.
- 1 Inch breadth.
- 5 Inch length.

WEIGHT 7 Ounce [1 ounce = 28.395]


LOCATION - Left hypochondriac region

• Partly epigastrium

Up to $9 - 11^{TH}$ ribs.

Spleen is non-palpable.

Anatomy of Spleen



EXTERNAL FEATURES

1. 2 ENDS	2. 3 BORDERS	3. 2 SURFACES
-----------	--------------	---------------

4. HILUM

- The ends are divided into two parts
- Anterior
- Posterior

SD ANATOMY NOTES

The borders are divided into three parts

- Superior
- Inferior
- Intermediate

The surface is further divided into two parts

- Diaphragmatic
- Visceral

ANTERIOR END

- Directed downwards.
- Laterally forward.

POSTERIOR END

- Directed upwards.
- Medially backward.

SUPERIOR BORDERS


- Near the anterior end.
- Intermediate border are round.

SURFACES

- Convex and smooth in nature.
- Related to the diaphragm.
- Visceral surface is concave and irregular.

LIGAMENTS

- Gastro splenic ligament.
- Lien renal ligament.
- Phrenic colic ligament.
- Phrenico splenic ligament.



PERITONEAL RELATION

- Gastro splenic ligament [hilum to spleen]
- Lien renal ligament [hilum to spleen]
- Phrenic colic ligament [supports anterior ends]

VISCERAL REALTIONS

- Related to fundus of stomach
- Anterior surface of left kidney
- Splenic flexure of colon
- Tail of pancreas

Gastric impression

Renal impression

Pancreatic impression

Hilum

Colic impression

Diaphragmatic surface - related to diaphragm

- Which seprated the spleen from pleura & lungs

FUNCTIONS

Phagocytosis.

- Hemiparesis.
- Immune response.
- Storage of red blood cells.

Blood Supply -

Splenic artery.

Nerve Supply -


Coeliac plexus - symphatic fibers

Venous supply - Drained by splenic vein

Tributaries are -

- 1. Short gastric
- 2. Left gastroepiploic vein
- 3. Pancreatic vein
- 4. Inferior mesenteric vein

Spleen Visceral and Diaphragmatic Surfaces



SUBCLAVIAN ARTERY

M56BOOKSTORE

Principal artery ehich continues as axillary artery

Supplies – part of neck & brain

Origin -

Right side – branch of brachio cephalic artery

Left side – it is branch of arch of aorta

Course -

Each artery arches lateraly from the sternoclvicular joint.

To outer border of 1st rib & continues with axillary artery

Scalenus anterior muscles crosses the artery Divides into 03 parts

Relations – SD NOTES

1st part -

Anterior – common carotid artery

- Vagus
- Internal jugular veins
- Sterno thyroid
- Sterno hyoid muscles
- Sternocleio mastoid

Posterior –

- supra pleural membrane
- cervical pleura
- apex of lung

2nd part –

Anterior – scalenus anterior

- Sternocleido mastoid

Posterior – cervical pleura

- Apex of lung

Superior – upper & middle trunks Of branchial plexus

3rd part –

Anterior – middle 1/3rd of clavicle

- Posterior border of sterno cleido mastoid

Posterior –

- Scalenus medius
- Brachial plexus
- Cervical pleura
- Apex of lung

Superior –

SD NOTES

- Upper & middle trunks of brachial plexus

Branches -

- Vertebral artery
- Internal thoracic artery
- Thyro cervical trunk
- Costo cervical trunk




Fig. 8.16: Origin and course of the subclavian arteries




Fig. 8.17: Course of subclavian and carotid arteries




Fig. 8.19: Branches of the subclavian artery. Note that the branches actually arise at different levels, but are shown at same level schematically




Fig. 8.20: Branches of the subclavian artery

SD NOTES M56BOOKSTORE

Thyroid gland

Thyroid gland – largest endocrine gland

- Lower part - in front

- Side of neck

Functions

- basal metabolic rate

- Calcium metabolism

- Natural iodine synthesis

Lobes -

1. Right lobe joined together by isthmus

2. Left lobe

Situation

- C5 – C7 & T1 vertebrae

- 5th tracheal ring

- Extend upto 2nd & 4th tracheal ring

Capsule

- true capsule

- False capsule

True capsule -

peripheral condensation of connective tissue of gland.

false capsule -

derived from – pretracheal layer of deep cervical fasica suspensory ligament of berry

PARTS - s.d notes

- Apex
- Base

SURFACES - 03

Lateral

Medial

Posteriolateral

Borders – 02

Anterior

Posterior

Isthmus – 02 surfaces -----anterior

Posterior

02 border ----- superior

Inferior

```
Relations -
```

Apex – sternothyroid muscle
Superior thyroid artery

External laryngeal artery

Base – inferior thyroid artery

Recurrent laryngeal nerve

Surface -

Lateral – sternothyroid

Strerno hyoid

Superior belly of omohyoid

Sternocleido mastoid

Medial – tracheal tube

Oesophagus

Inferior constriction

Crico thyroid

Recurrent laryngeal nerve

Postero lateral – carotid sheath


Border – superior thyroid artery
Inferior thyroid artery
Parathyroid gland
Thoracic duct


Arterial supply –
Superior thyroid artery
Inferior thyroid artery
Accessory thyroid artery

Venous drainage –
Superior thyroid vein
Middle thyroid vein
Inferior thyroid vein


Lymphatic – deep cervical lymph nodes .

B. RELATIONS







a.Arterial supply



b.Venous drainage



D. APPLIED



Reference ----- Anatomy – $bdc - 9^{th}$ edition – 160- 3^{rd} volume .

TONGUE ANATOMY

M56BOOKSTORE----DR.SD

MUSCULAR ORAGN

SITUTATED ---Floor of mouth

Functions ----

- 1. Taste
- 2. Speech
- 3. Chewing
- 4. Deglution

Structure -----

- 1. Root
- 2. Tip
- 3. Body

Tip of the tongue -----

- Anterior free end
- Behind upper incisor teeth

Root of the tongue -----

- Attached to styloid process and soft palate above
- To mandible and hyoid bone

Body of the tongue -----

- 02 borders
- 02 surfaces

-

DR SD ---- M56BOOKSTORE

- 02 borders -----
- Right lateral borders
- Left lateral borders
- 02 surfaces -----
- Dorsal surfaces / dorsum of tongue
- Ventral surfaces
- 1. Dorsal surafces of tongue -----
- 1. Oral part
- 2. Pharyngeal part

Convex antero posteriotly

- 1. Anterior 2/3rd or oral part
- 2. Posterior 1/3rd or pharyngeal part

Anterior 2/3rd & post 1/3rd -----

Joined by sulcus terminalis (v – shaped)

Papillae are present on anterior 2/3rd portion of tongue ----

Papillae are projections of mucous membrane

Rough ness present on anterior 2/3rd part

DR SD ---- M56BOOKSTORE

Papillae types ----

- 1. Cicrcumvallate papillae -----lateral wall
- It is of v shape
- 1-2 mm in diameter
- 8-12 in number
- Fungiform papillae --- Smaller and more in number
 Large in size than filliform papillae
- 3. Fillifom papillae ---
- Present in the middle part of the dorsum tongue
- Smallest in size
- 4. Foliate papillae -----
- 3-4 small vertical folds on the margin of sulcus terminalis

Posterior 1/3rd part -----

- Lies behind the palatoglossal arches and sulcus terminalis
- Part (lymphoid follicles are present)
- =
- Collectivey called lingual tonsli
- Mucous glands are present
- Either side of median fold ----depression called valleculla

Muscles of tongue -----

- 1. Intrinsic muscle
- 2. Extrinsic muscle
- Intrinsic muscles ----
 - 1. Superior longitudnal linguae ----concave shortness
 - 2. Inferior longitudnal linguae -----convex shorten
 - 3. Transverse linguae -----narrow & elongated
 - 4. Verticalis linguae ----- broad and flattened

DR SD ---- M56BOOKSTORE

Extrinsic muscles ----

- 1. Genioglossus -----protudes
- 2. Hyoglossus ----- depresses
- 3. Styloglossus ----rectracts
- 4. Palatoglossus----- elevates

Arterial supply -----

- lingual artery
- branched of carotid artery

venous drainage ----

deep lingual vein

Nerve supply -----

All muscle ----hypoglossal nerve ----12th cranial)

Motor nerve -----except -----palatoglossus ----paryngeal plexus

Sensory ----- Anterior 2/3rd ---lingual nerve , posterior1/3rd ---glossopharybgeal nerve – 9th CN

CLINICAL ANATOMY -----

HYPOGLOSSAL nerve injury -----paralysis of tongue muscles

Glossitis -----generalized ulceration

Stomatitis ---- inflammation of tongue

Carcinoma of tongue

Pale colour in anemia


Violet colour in riboflavin defeciency

Dry in dehydration


Blue in cyanosis


Tongue bite -----

TASTE PATHWAY



OF THE WARREN





BREAKDOWN OF TONGUE DEVELOPMENT -----

• Embryonic origin:

- The tongue begins forming around the 4th week of embryonic development.
- It develops from multiple pharyngeal arches (first, second, and third).

• Formation of the anterior 2/3 (body):

- The anterior two-thirds of the tongue originates from the **first pharyngeal arch**.
- It starts as **two lateral lingual swellings** and a **tuberculum impar**. The lateral swellings grow and fuse, covering the tuberculum impar to form the anterior portion.
- The epithelium of this region is innervated by the **lingual nerve** (branch of the mandibular nerve, CN V3).

• Formation of the posterior 1/3 (root):

- The posterior one-third of the tongue forms from the **third pharyngeal arch**, with contributions from the second and third arches.
- This portion develops from the **hypobranchial eminence**.
- The mucosa of the posterior part is innervated by the **glossopharyngeal nerve** (CN IX).

• Development of the muscles:

- The muscles of the tongue originate from **occipital myotomes**, specifically from somites (segments of mesoderm).
- These muscles are innervated by the **hypoglossal nerve** (CN XII), which is responsible for tongue movement.

• Development of taste buds:

- The taste buds develop from interactions between epithelial and mesenchymal tissues.
- The anterior two-thirds (innervated by the **chorda tympani**, a branch of the facial nerve CN VII) provides taste sensation.
- The posterior one-third (innervated by the **glossopharyngeal nerve** CN IX) is also responsible for taste.

• Fusion and appearance of the median sulcus:

• The fusion of the lateral lingual swellings creates a midline structure called the **median** sulcus.


• Developmental milestones:


• By around week 8 of embryonic development, the tongue is largely formed.

• Functional development and maturation of taste sensation continue into the fetal stage.

• Innervation summary:

- Motor: Hypoglossal nerve (CN XII) for movement.
- **General sensory**: Lingual nerve (CN V3) for anterior 2/3, glossopharyngeal nerve (CN IX) for posterior 1/3.
- **Taste sensory**: Chorda tympani (CN VII) for anterior 2/3, glossopharyngeal nerve (CN IX) for posterior 1/3.





Urinary bladder

M56bookstore

- > Intrapelvic organ
- > Storehouse of urine
- > Tetrahedral shape empty conition
- ➤ Ovoid shape filled condition

External features -

- > Apex
- Base
- ➤ Neck
- Surfaces
- ➤ Borders
- ➤ Apex directed downwards in bladder
- ➤ Base directed backwards in empty bladder
- ➤ Neck most fixed part urethral sphincter started

Surfaces - 03 surfaces -

- > Superior surface
- ➤ Right surface
- ➤ Left inferolateral surface

Borders – 04 borders

- > 02 lateral
- > 01 anterior
- > 01 posterior


Capacities -


- ➤ Average capacity 120-320 ml
- ➤ Mean capacity 220 ml
- ➤ Sense of filling 100 150 ml
- ➤ Micturition 250 ml
- Painful sensation >450 ml

Male urethra parts -

- > Pre postatic
- > Prostatic part
- > Membraneous part
- important parts
- > Spongy part of urethra
- > Penile part
- > External urethral orifice

Diagram - image based -





Relations - Males -

- ➤ Apex median umblical ligament
- ➤ Base Ampulla of rectum
- Seprated byRectovesical pouch

> Pair of seminal vesicles

- > Vas deferens
- > Recto vesical pouch
- > Fasica of denon Villiers

Females –

- > Vagina
- > Supra vaginal part of cervix

Superior surface – realation

- > Triangular
- > Covered with peritoneum
- ➤ Uterus

Inferiolateral surface – Relations

- Directed downwards & laterally
- Pelvic surface
- > Fasicia covering levator ani & obturator internus muscles
- > Sapace of retzius

Space of retzius - 2 marks -

> Space b/w infero lateral urinary bladder & fascia covering levator ani .

Neck Relations –

- Lowest part of urinary bladder
- Urethra begins male prostatic part includes
- Sourrounded base of prostate

Females -

> contact with urogenital diaphragm


Distended bladder / empty -

- > Shape ovoid
- ➤ Surfaces –
- ➤ anterior inferior No peritoneum
- ➤ Posterior superior with peritoneum
- > 02 lateral surfaces

Parts -

> summit , base , neck

Examination of bladder wall – tube of cystoscope – (imp)



Ligaments -

- > True ligaments
- > False ligaments

True ligaments -

- ➤ Median umblical ligament
- ➤ Medial puboprostatic ligament
- ➤ Lateral puboprostatic ligament
- > Lateral ligament pair
- ➤ Posterior ligament pair

False ligament –

- Median umblical fold
- > Medial umblical fold
- > Lateral false ligament
- > Posterior false ligament

Structure -

- Outside to inside –
- > Serous
- > Muscularis -----detrusor muscle
- > Submucosa
- Mucosa

Trigonum vesicae –

> Ureteric ridge of mercier

Uvula vesicae –

Produced by median lobe of prostate

Arterial supply -

- Superior vesicle artery
- > Inferior vesicle artery
- Obturator artery
- > Branch of inf gluteal artery
- Female uterine artery

Venous drainage -

➤ Venis form -----space of retzius -----communicates with prostatic plexus -----finally ------internal iliac veins .

Lymph nodes -

- > External iliac nodes
- > Internal iliac nodes

Nerve supply –

- ➤ Parasympathetic splanchic nerve -----s2 s3 s4
- ➤ Sympathic plexus hypogastric fibers ----T11 L2
- > Somatic pudendal nerve

Applied anatomy –


- > Hypertrophy of bladder
- > Urinary bladder incontinence
- > Trabeculated bladder
- Dribbling of urine through autonomus bladder
- ➤ Bladder rupter any injury of lower abdominal wall

Suprapubic cystotomy (spc)


- > Pt. cannot empty the bladder due to urethral trauma or bony fracture.
- ➤ Bladder stones --- concentrated urine.

Diagrams –


A. PARTS




B. LIGAMENTS




C. RELATIONS IN MALE



D. RELATIONS IN FEMALE






M56bookstore

VENTRICLES OF THE BRAIN - M56BOOKSTORE

VENTRICLES OF THE BRAIN -

COMMUNICATIONG CAVITY WITHIN THE CNS THAT ARE LINED BY EPENDYMAL CELLS, WHICH PRODUCES, CONTAINS AND CIRCULATES CSF.



CAVITIES -

VENTRICULAT SYSTEM COMPRISES –

TWO LATERAL VENTRICALS


3RD VENTRICAL

4TH VENTRICAL

TERMINAL VENTRICAL

THESE CHANNELS ARE -

INTERVENTRICULAR FORAMEN OR FORAMEN OF MONRO
CEREBRAL AQUEDUCT
CENTRAL CANAL



VENTRICLES -

LATERAL VENTRICAL –

BODY - LIES IN THE PARIENTAL LOBE

ROOF – CORPUS CALLOSUM

FLOOR – BODY OF CAUDATE NUCLEUS


UPPER SURFACE OOF THALAMUS


CHOROID PLEXUS

BODY OF FORNIX

MEDIAL WALL - SEPTUM PELLUCIDUM

LATERAL WALL – NARROW AREA AT ROOF AND FLOOR





ANTERIOR HORN -

LOCATION - FORNTAL LOBE


ROOF - CORPUS CALLOSUM

FLOOR - CORPUS CALLOSUM,

CAUDATE NUCLEUS

ANTERIOR – CORPUS CALLOSUM

MEDIALLY - SEPTUM PELLUCIDUM



POSTERIOR HORN -

LOCATION OCCIPITAL LOBE

ROOF & LATERAL WALL – TAPETUM OF CORPUS CALLOSUM

MEDIALLY – 2 ELEVATIONS – BULB OF POSTERIOR HORN, CALCAR AVIS


PRODUCE BY CALCARINE SULCUS.

INFRIOR HORN –

LOCATION - TEMPORAL LOBE

ROOF - TAPETUM, TAIL OF CAUDATE NUCLEUS, AMYGDALOID NUCLEUS, STRIA TERINALIS

FLOOR - HIPPOCAMPUS, FIMBRIA, HIPPOCAMPUS AND COLLATERAL EMINENCE.



3RD VENTRICLE -

LOCATION – B/W TWO THALAMUS


ROOF - FORNIX, TELLA CHOROIDA

FLOOR - HYPOTHALAMUS

LATERALLY – THALAMUS & HYPOTHALAMUS

ANTERIORLY – LAMINA TERMINALIS

POSTERIORLY – OPENING INTO THE CEREBRAL AQUEDUCT, PINEAL RECESS.



4TH VENTRICLE -

LOCATION – ANTERIOR – CEREBELLUM

POSTERIOR – PONS


SUPERIOR ½ OF MEDULLA OBLANGATA


FLOOR – POST. SURFACE OF PONS

CRANIAL ½ OF MEDULLA OBLONGATA

LATERALLY – INF. CEREBELLAR PEDUNCLE & SUP PEDUNCLE

ROOF – TENT SHAPED ROOF INTO CEREBELLUM





FUNCTIONS OF VENTRICLES -

CSF – WATER LIKE FLUID RESIDES WITHIN THE VENTRICULAR SYSTEM.


Appearance and composition


Appearance	Clear and colorless
Volume	150 mL
Rate of production	0.5 mL/minute
Pressure (spinal tap with patient in lateral recumbent position)	60-150 mm of water
Composition	
Protein	15-45 mg/100 mL
Glucose	50-85 mg/100 mL
Chloride	720-750 mg/100 mL
Number of cells	0-3 lymphocytes/cumm

CSF FORMATION -

CHOROID PLEXUS – IT IS A VASCULAR FRINGE COMPOSED OF PIAMATER COVERED WITH THE EPENDYMAL LINING OF VENTRICULAR CAVITY


CIRCULATION OF CSF -





FACTORS OF CSF FLOW -

PULSATION OF CEREBRAL & SPINAL ARTERIES
MOVEMENT OF VERTEBRAL COLUMN
RESPIRATION AND COUGHING
CHANGING OF POSITION.




Function of the CSF

- Cushions and protects the central nervous system from trauma
- Provides mechanical buoyancy and support for the brain
- Serves as a reservoir and assists in the regulation of the contents of the skull
- Nourishes the central nervous system
- Removes metabolites from the central nervous system
- Serves as a pathway for pineal secretions to reach the pituitary gland

APPLIED ANATOMY -


Papilledema is a term that is exclusively used when a disc swelling is secondary to increased intracranial pressure (ICP).





Hydrocephalus -

It is the buildup of fluid in the cavities (ventricles) deep within the brain. The excess fluid increases the size of the ventricles and puts pressure on the brain.



Causes can be:

- 1. Blockage of the circulation of cerebrospinal fluid
- 2. Diminished absorption of cerebrospinal fluid
- 3. Excessive formation of cerebrospinal fluid

Diseases involving CFS

- Meningitis
- Blockage of the subarachnoid space in the vertebral canal
- Tumors of the fourth ventricle

